

2000 Ambient Air Quality Report

STATE OF NORTH CAROLINA

Michael F. Easley, Governor

DEPARTMENT OF

ENVIRONMENT

AND

NATURAL RESOURCES

William G. Ross, Jr., Secretary

DIVISION OF AIR QUALITY

B. Keith Overcash, P.E., Director

AMBIENT MONITORING SECTION

Hoke P. Kimball, Chief

PUBLISHED

December 2002

2000 Ambient Air Quality Report

Ambient Monitoring Section Report # 2002.01

Acknowledgements:

Primary Author: Neil Berg
Additional Assistance: G. Stephen Few

Public Sources of Data:

North Carolina
Division of Air Quality <http://daq.state.nc.us/>

Forsyth County
Environmental Affairs
Department <http://www.co.forsyth.nc.us/EnvAffairs/weathereport.htm>

EPA/AIRS Air Quality
Subsystem <http://www.epa.gov/airsweb>

National
Atmospheric
Deposition Program <http://nadp.sws.uiuc.edu>

Published: December 2002

Not copyrighted.

200 copies of this public document were printed on recycled paper at a total cost of \$518.72, or \$2.59 per copy.

This document is printed on recycled paper.

Preface

This report is issued by the Division of Air Quality of the Department of Environment and Natural Resources to inform the public of air pollution levels throughout the state of North Carolina. It describes the sources and effects of the following pollutants for which the U.S. Environmental Protection Agency and the State of North Carolina have established ambient air quality standards:

Particulate Matter
Carbon Monoxide
Sulfur Dioxide

Nitrogen Dioxide
Ozone
Lead

A brief discussion of the ambient air monitoring program, including a description of the monitoring network, is provided. Detailed results are presented of monitoring that was conducted in 2000 to measure the outdoor concentrations. The data are presented graphically and as statistical summaries, including comparisons to the ambient air quality standards. The report discusses the recorded data, seasonal variability of some pollutants. Data and areas exceeding the ambient air quality standards are identified. Factors that have contributed to those exceedances are described also.

Acid rain data from the National Atmospheric Deposition Program/National Trends Network for North Carolina also are included for 2000. Data collected after 2000 will be discussed in later reports.

Current air pollution information is available to the public 24 hours a day through the use of the air quality index telephone numbers listed below:

Statewide toll-free
(for Asheville, Durham, Fayetteville, Greensboro, Greenville,
Raleigh, Wilmington, and Winston-Salem areas)

888-AIR-WISE

Charlotte area

704-333-SMOG

Additional copies of this report and previous annual reports are available from:

Division of Air Quality
Department of Environment and Natural Resources
1641 Mail Service Center
Raleigh, North Carolina 27699-1641

Comments regarding this report or suggestions for improving future reports are welcomed. Comments may be sent to Dr. Wayne L. Cornelius, at the above address.

B. Keith Overcash, P.E., Director
Division of Air Quality

Executive Summary

In 2000, the North Carolina Division of Air Quality (DAQ), the three local program agencies and one tribal agency (listed in Appendix A) collected 485,655 air quality samples. These samples included measurements of the U.S. Environmental Protection Agency's (EPA) criteria air pollutants: particulate matter, carbon monoxide, ozone, sulfur dioxide, nitrogen dioxide and lead. This report discusses each pollutant and presents summary tables, maps, charts and explanations of the data.

The report also includes data from weekly acid rain samples collected by the National Atmospheric Deposition Program/National Trends Network (NADP) at seven North Carolina sites and one Tennessee site very close to the North Carolina border. It discusses acid rain and presents summary tables, maps, charts and explanations of the data.

Three different types of **particulate matter** were sampled in North Carolina during 2000. Total Suspended Particulate (TSP), considered to be particles having an aerodynamic diameter of 100 micrometers or less, is regulated by North Carolina standards. Particulate matter (PM₁₀) with a mean aerodynamic diameter less than or equal to a nominal 10 micrometers (0.00004 inches) is regulated by both EPA and N.C. standards. Fine particulate matter (PM_{2.5}) with a mean aerodynamic diameter less than or equal to a nominal 2.5 micrometers (0.00001 inches) has been regulated by EPA since 1997.

TSP was sampled at 2 sites, yielding 92 daily samples. No exceedances of the state TSP ambient air quality standard for 24-hour samples (150 µg/m³) were observed in 2000.

PM₁₀ was sampled at 32 sites, yielding 2628 daily samples. There were no exceedances of the National Ambient Air Quality Standards for PM₁₀ (150 µg/m³ for 24-hour samples and 50 µg/m³ for the annual arithmetic mean). Mean 24-hour concentrations have decreased about 30 percent since 1985.

PM_{2.5} was sampled at 38 sites yielding 5188 daily samples. There was one exceedance of the ambient air quality standards for PM_{2.5} (65 µg/m³ for 24-hour samples). The value of 67.4 was observed at the Raeford Road site in Fayetteville. Twenty one-of the 38 sites exceeded the annual arithmetic mean standard of 15 µg/m³.

Carbon monoxide (CO), the most common air pollutant, largely results from fuel combustion. The most likely areas to have excessive CO concentrations are larger cities where there are more cars and congested streets.

CO was sampled at 12 sites, yielding 61,959 valid hourly averages. The National Ambient Air Quality Standards for CO are 35 ppm for the maximum one-hour average and 9 ppm for the maximum eight-hour average. There were no exceedances of the standards. The highest one-hour concentration of 14.1 was observed at the Tryon site in Charlotte. The highest eight hour concentration of 5.4 ppm was observed at the Oleander and College site in Wilmington and the

Person Street site in Raleigh. Both the mean one-hour average and the mean eight-hour average have been decreasing by about 4 percent per year. The combined effects of newer cars in the vehicle fleet, traffic control strategies, and the Inspection and Maintenance program in Durham, Orange, Wake, Forsyth, Guilford, Cabarrus, Gaston, Mecklenburg, and Union Counties have helped reduce the number and intensity of CO exceedances from previous years.

Ozone (O₃) forms in the lower atmosphere when hydrocarbons (or volatile organic compounds) and nitrogen oxides chemically react in the presence of sunlight and high temperatures. The main emphases in control of ozone has been to limit hydrocarbon and nitrogen oxide emissions.

O₃ was sampled at 45 sites, yielding 226,922 valid hourly averages. The National Ambient Air Quality Standard for O₃ is 0.08 ppm for the maximum eight-hour average and 0.12 ppm for the maximum one-hour average.

In 2000, there were 6 exceedances of the one-hour standard, all of which occurred on three days in June. Twenty exceedances occurred in North Carolina in 1999, and 24 occurred in 1998. Mecklenburg, Rowan and Wake Counties met or exceeded the criteria for nonattainment of the one-hour ozone standard with nine, five and five exceedances respectively over a three-year period, however EPA had rescinded the one-hour standard during that time period. Mecklenburg County was redesignated as in attainment for ozone in July 1995. Hydrocarbon control strategies continue to be used there to help reduce ozone concentrations.

In 2000, the 8-hour standard was exceeded 239 times, on 35 different days, with five counties having 10 or more exceedances at individual sites. The site at 246 Main Street, Cooleeemee in Davie County had the highest number, 17.

Sulfur dioxide (SO₂) is mainly produced by combustion of fossil fuels containing sulfur compounds and the manufacture of sulfuric acid.

SO₂ was sampled at 24 sites, yielding 176,178 valid hourly averages. There were no exceedances of the National Ambient Air Quality Standards (365 µg/m³ or 0.14 ppm for a 24-hour average, 1300 µg/m³ or 0.50 ppm for a three-hour average, 80 µg/m³ or 0.03 ppm for the annual arithmetic mean) at network monitoring sites.

Nitrogen oxides (NO_x) are produced primarily from the burning of fossil fuels such as coal, oil and gasoline, due to the oxidation of atmospheric nitrogen and nitrogen compounds in the fuel. The primary combustion product is NO, which reacts with hydrocarbons, ozone and other atmospheric compounds to form NO₂. NO_x compounds play an important role in the formation of ozone. Reactive nitrogen species (NO_y) were monitored in Charlotte, Raleigh, and Winston-Salem to gather data for the development of control strategies for ozone non-attainment areas.

The criteria pollutant NO₂ was sampled at two sites, yielding 12,688 valid hourly averages. There were no exceedances of the National Ambient Air Quality Standard (0.053 ppm for the

annual arithmetic mean). The mean one-hour average concentration has been decreasing by about 1.5 percent per year.

Lead (Pb) emissions result from coal combustion and the sandblasting of highway bridges, overpasses, and water tanks. In the past, the combustion of gasoline containing tetraethyl lead as an additive was a major source.

Lead was not sampled in 2000. There have been no recent exceedances of the ambient air quality standard for lead ($1.5 \mu\text{g}/\text{m}^3$ for a quarterly arithmetic mean). Mean lead concentrations have decreased by 92 percent since 1979. The steady decline in the use of leaded gasoline is primarily responsible for this trend.

Acid Rain is produced when nitrate and sulfate ions from motor vehicles, combustion and industrial sources reach the upper atmosphere, react with moisture in the air, and are deposited as acid precipitation. Monitoring of pH and other ion concentrations in precipitation will help to identify trends and demonstrate the results of efforts to reduce emissions from mobile and industrial sources.

The annual mean pH in 2000 ranged from 4.39 (Rowan County) to 4.73 (Sampson County).

CONTENTS

1	Introduction.....	1
2	Description of Criteria Pollutants	2
2.1	Particulate Matter.....	2
2.1.1	Sources.....	2
2.1.2	Effects	3
2.2	Carbon Monoxide	3
2.2.1	Sources.....	3
2.2.2	Effects	3
2.3	Ozone	4
2.3.1	Sources.....	4
2.3.2	Effects	4
2.4	Sulfur Dioxide	4
2.4.1	Sources.....	4
2.4.2	Effects	5
2.5	Nitrogen Oxides	5
2.5.1	Sources.....	5
2.5.2	Effects	5
2.6	Lead	6
2.6.1	Sources.....	6
2.6.2	Effects	6
3	Standards.....	7
4	Ambient Air Quality Monitoring Program.....	9
5	Pollutant Monitoring Results	17
5.1	Total Suspended Particulates	17
5.2	PM ₁₀	20
5.3	PM _{2.5}	24
5.4	Carbon Monoxide	28
5.5	Ozone	32
5.6	Sulfur Dioxide	44
5.7	Nitrogen Dioxide	48
5.8	Lead	50
6	Air Quality Index.....	51
7	Acid Rain	56
7.1	Sources.....	56
7.2	Effects	56
7.3	Monitoring	56
	References.....	60
	Appendix A. Air Pollution Monitoring Agencies.....	61
	Appendix B. Exceptional Events	64
	Appendix C. Box-And-Whisker Plots	65
	Appendix D. Nonattainment and North Carolina	66

List of Tables

Table 3.1 National and North Carolina Ambient Air Quality Standards	8
Table 4.1 Ambient Air Monitoring Sites Operated in North Carolina, 2000	11
Table 5.1 Total Suspended Particulates in Micrograms Per Cubic Meter for 2000	19
Table 5.2 PM ₁₀ in Micrograms Per Cubic Meter for 2000.....	21
Table 5.3 PM _{2.5} in Micrograms Per Cubic Meter for 2000	25
Table 5.4 Carbon Monoxide in Parts Per Million for 2000	30
Table 5.5 One-Hour Ozone in Parts Per Million for 2000	34
Table 5.6 Eight-Hour Ozone in Parts Per Million for 2000	37
Table 5.7 Sulfur Dioxide in Parts Per Million from All Sites for 1998-2000	45
Table 5.8 Nitrogen Dioxide in Parts Per Million for 2000	49
Table 7.1 pH, Conductivity in Microsiemens per Centimeter and Precipitation in Inches from the National Atmospheric Deposition Program for 2000.....	58
Table 7.2 Ion Concentrations in Milligrams per Liter (Precipitation-weighted Annual Means) from the National Atmospheric Deposition Program Data for 2000.....	59

List of Figures

Figure 4.1 Monitoring Sites Active in 2000	10
Figure 5.1 Location of PM ₁₀ Monitoring Sites.....	20
Figure 5.2 PM ₁₀ : Second Highest 24-Hour Averages, 2000	23
Figure 5.3 PM ₁₀ : Maximum Annual Arithmetic Means, 2000	23
Figure 5.4 Location of PM _{2.5} Monitoring Sites	24
Figure 5.5 PM _{2.5} : Second Highest 24-Hour Averages, 2000	27
Figure 5.6 PM _{2.5} : Maximum Annual Arithmetic Means, 2000	27
Figure 5.7 Location of Carbon Monoxide Monitoring Sites	29
Figure 5.8 Carbon Monoxide: Second Highest 1-Hour Average, 2000	31
Figure 5.9 Carbon Monoxide: Second Highest Non-overlapping 8-Hour Average, 2000	31
Figure 5.10 Location of Ozone Monitoring Sites	33
Figure 5.11 Ozone: Second Highest Annual 1-Hour Average, 2000	40
Figure 5.12 Ozone: Second Highest Annual 8-Hour Average, 2000	40
Figure 5.13 Distribution of Statewide Fourth-Maximum 8-Hour Ozone Concentrations, 1991- 2000, and Smoothed Regression Trend Line.....	41
Figure 5.14 Number of Exceedances of the Daily Maximum 8-Hour Ozone Averages of 0.085 ppm or Greater, 1993-2000.....	42
Figure 5.15 Number of Exceedances of the Daily Maximum 8-Hour Ozone Averages of 0.085 ppm or Greater, by Region, 1993-2000.	43
Figure 5.16 Location of Sulfur Dioxide Monitoring Sites	44
Figure 5.17 Sulfur Dioxide: Second Highest 3-Hour Averages in the Most Recent Year of Data from 1998, 1999 or 2000	47
Figure 5.18 Sulfur Dioxide: Second Highest 24-Hour Averages in the Most Recent Year of Data from 1998, 1999 or 2000	47
Figure 5.19 Location of Nitrogen Dioxide Monitoring Sites	48
Figure 6.1 Daily Air Quality Index Values for Asheville, NC, Metropolitan Statistical Area, 2000	53
Figure 6.2 Daily Air Quality Values for Charlotte-Gastonia, NC,-Rock Hill, SC, Metropolitan Statistical Area, 2000	53
Figure 6.3 Daily Air Quality Index Values for Fayetteville, NC, Metropolitan Statistical Area, 2000	54
Figure 6.4 Daily Air Quality Index Values Greensboro- Winston-Salem-High Point, NC, Metropolitan Statistical Area, 2000	54
Figure 6.5 Daily Air Quality Index Values for Raleigh-Durham, NC, Metropolitan Statistical Area, 2000.....	55
Figure 6.6 Daily Air Quality Index Values for Wilmington, NC, Metropolitan Statistical Area, 2000	55
Figure 7.1 Annual Mean pH Values at North Carolina NADP Sites, 2000	57

1 Introduction

This annual report summarizes the ambient air monitoring performed in calendar year 2000 by the North Carolina Division of Air Quality (DAQ), three local air pollution agencies and one tribal agency, which are more fully described in Appendix A.

There were 485,655 air quality samples of the U.S. Environmental Protection Agency's (EPA) criteria pollutants B particulate matter, carbon monoxide, ozone, sulfur dioxide, nitrogen dioxide and lead - which are discussed in this report.

Chapter 2 describes the criteria pollutants and discusses their sources and effects on human health, plants and animals. Chapter 3 outlines the standards applied to criteria pollutant concentrations established by the EPA and the state of North Carolina to protect human health (primary standards) and plants, animals, and property (secondary standards). Chapter 4 describes the ambient monitoring program conducted by DAQ and three local program agencies. Chapter 5

gives detailed monitoring results for each pollutant, with a map of the monitor sites, a table of the monitor summary statistics relevant to the standards, one or more maps summarizing the important statistics for each county with monitors, and additional summaries as appropriate to each pollutant. Chapter 6 describes the EPA Air Quality Index for the criteria pollutants and charts index measurements for five Metropolitan Statistical Areas of North Carolina. Chapter 7 presents sources, effects and monitoring of acid rain data conducted in North Carolina by the National Atmospheric Deposition Program and National Trends Network (NADP). It also includes a map of the calendar year mean pH level and site statistics for the calendar year in two tables.

In previous reports, we included a final chapter that provides statewide summary trends for the criteria pollutants. However, due to the budget shortfall, the trends section of this report will be omitted. A stand alone trends report will be prepared at a later date.

2 Description of Criteria Pollutants

2.1 Particulate Matter

Atmospheric particulate matter is defined as any airborne material, except uncombined water (liquid, mist, steam, etc.) that exists in a finely divided form as a liquid or solid at standard temperature (25°C) and pressure (760 mm mercury) and has an aerodynamic diameter of less than 100 micrometers. In the period covered by this report, three sizes of particulate matter were monitored, total suspended particulate (TSP), PM₁₀ and PM_{2.5}. TSP is any particulate matter measured by the method described in EPA regulations 40 CFR 50 App. B (United States Environmental Protection Agency [US EPA] 1993, p. 715-728) and is generally considered to be particles having an aerodynamic diameter of 40 µm or less (Watson and Chow 2001), although particles up to about 100 µm are sometimes captured by samplers. (The probability of inhalation for 100 µm particles is about 50 percent and increases with decreasing particle size [Maynard and Jensen 2001].) PM₁₀ is particulate matter with an aerodynamic diameter less than or equal to 10 µm as measured according to EPA regulations 40 CFR 50 App. J (United States Environmental Protection Agency [US EPA] 1993, p. 769-773). TSP measurements have

been made in North Carolina since the early 1960s and PM₁₀ has been sampled locally in Charlotte since 1985 and statewide since 1986 (North Carolina Department of Environment, Health, and Natural Resources 1991). The new PM_{2.5} standard was adopted by North Carolina on April 1, 1999. On May 14, 1999 the U.S. Court of Appeals ruled the setting of the standard by EPA was an unconstitutional use of authority and could be vacated. The Supreme Court later upheld the new standard. EPA continues to require monitoring for PM_{2.5}.

2.1.1 Sources

Particulates are emitted by many human activities, such as fuel combustion, motor vehicle operation, industrial processes, grass mowing, agricultural tilling and open burning. Natural sources include windblown dust, forest fires, volcanic eruptions, and plant pollen.

Particles emitted directly from a source may be either fine (less than 2.5 µm) or larger (2.5 - 60 µm), but particles formed in the atmosphere will usually be fine. Generally, larger particles have very slow settling velocities and are characterized as suspended particulate matter. Typically, fine particles originate by condensation of

materials produced during combustion or atmospheric reactions.

2.1.2 Effects

Particulate matter can cause health problems affecting the breathing system, including aggravation of existing lung and heart disease, limitation of lung clearance, changes in form and structure of organs, and development of cancer. Individuals most sensitive to the effects of particulate matter include those with chronic obstructive lung or heart disease, those suffering from the flu, asthmatics, the elderly, children, and mouth breathers.

Health effects from inhaled particles are influenced by the depth of penetration of the particles into the respiratory system, the amount of particles deposited in the respiratory system, and by the biological reaction to the deposited particles. The risks of adverse health effects are greater when particles enter the tracheobronchial and alveolar portions of the respiratory system. Small particles can penetrate into these deeper regions of the respiratory system. Healthy respiratory systems can trap particles larger than 10 micrometers more efficiently before they move deeply into the system and can more effectively remove the particles that are not trapped before deep movement.

Particulate matter also can interfere with plant photosynthesis, by forming a film on leaves reducing exposure to sunlight. Particles also can cause soiling and degradation of property, which can be costly to clean and maintain.

Suspended particles can absorb and scatter light, causing reduction of visibility. This is a national concern, especially in areas such

as national parks, historic sites and scenic attractions visited by sightseers.

2.2 Carbon Monoxide

Carbon monoxide (CO) is the most commonly occurring air pollutant. CO is a colorless and poisonous gas produced by incomplete burning of carbon-containing fuel.

2.2.1 Sources

Most atmospheric CO is produced by incomplete combustion of fuels used for vehicles, space heating, industrial processes and solid waste incineration. Transportation accounts for the majority of CO emissions. Boilers and other fuel burning heating systems are also significant sources.

2.2.2 Effects

Breathing carbon monoxide affects the oxygen-carrying capacity of the blood. Hemoglobin in the blood binds with CO more readily than with oxygen, starving the body of vital oxygen.

Individuals with anemia, lung and heart diseases are particularly sensitive to CO effects. Low concentrations affect mental function, vision and alertness. High concentrations can cause fatigue, reduced work capacity and may adversely affect fetal development. Chronic exposure to CO at concentrations as low as 70 ppm (80 mg/m³) can cause cardiac damage. Other health effects associated with exposure to CO include central nervous system effects and pulmonary function difficulties.

Ambient CO apparently does not adversely affect vegetation or materials.

2.3 Ozone

Ozone is a clear gas that forms in the troposphere (lower atmosphere) by chemical reactions involving hydrocarbons (or volatile organic compounds) and nitrogen oxides in the presence of sunlight and high temperatures. Even low concentrations of tropospheric ozone are harmful to people, animals, vegetation and materials. Ozone is the most widespread and serious criteria air pollutant in North Carolina.

Ozone in the upper atmosphere (stratosphere) shields the earth from harmful effects of ultraviolet solar radiation. Stratospheric ozone can be damaged by the emission of chlorofluoro-hydrocarbons (CFCs) such as Freon.

2.3.1 Sources

Ozone (O₃) is the major component of a complex mixture of compounds known as photochemical oxidants. Ozone is not usually emitted directly into the atmosphere, but is formed by a series of complex reactions involving hydrocarbons, nitrogen oxides and sunlight. Ozone concentrations are higher during the daytime in late spring, summer and early autumn when the temperature is above 60°F and the sunlight is more intense.

Two natural sources of upper atmosphere ozone are solar radiation and lightning during thunderstorms. These are not

significant sources of tropospheric (ground level) ozone.

2.3.2 Effects

Ozone is a pulmonary irritant, affecting the respiratory mucous membranes, as well as other lung tissues and respiratory functions. Ozone has been shown to impair normal function of the lung causing shallow, rapid breathing and a decrease in pulmonary function. Other symptoms of exposure include chest tightness, coughing and wheezing. People with asthma, bronchitis or emphysema probably will experience breathing difficulty when exposed to short-term concentrations between 0.15 and 0.25 ppm. Continued or repeated long-term exposure may result in permanent lung structure damage.

Ozone damages vegetation by injuring leaves. Ozone also accelerates material aging, cracking rubber, fading dyes and eroding paint.

2.4 Sulfur Dioxide

Sulfur dioxide (SO₂) is a colorless, corrosive, harmful gas with a pungent odor. Smaller concentrations of sulfur trioxide and other sulfate compounds are also found in SO₂ emissions. Sulfur oxides contribute to the formation of acid rain and the formation of particles that reduce visibility.

2.4.1 Sources

The main sources of SO₂ are combustion of fossil fuels containing sulfur compounds and the manufacture of sulfuric acid. Other

sources include refining of petroleum and smelting of ores that contain sulfur.

2.4.2 Effects

The most obvious health effect of sulfur dioxide is irritation and inflammation of body tissues brought in contact with the gas. Sulfur dioxide can increase the severity of existing respiratory diseases such as asthma, bronchitis, and emphysema. Sulfuric acid and fine particulate sulfates, which are formed from sulfur dioxide, also may cause significant health problems. Sulfur dioxide causes injury to many plants. A bleached appearance between the veins and margins on leaves indicates damage from SO_2 exposure. Commercially important plants sensitive to SO_2 include cotton, sweet potatoes, cucumber, alfalfa, tulips, apple trees, and several species of pine trees.

2.5 Nitrogen Oxides

Several gaseous oxides of nitrogen are normally found in the atmosphere, including nitrous oxide (N_2O), nitric oxide (NO) and nitrogen dioxide (NO_2). Nitrous oxide is a stable gas with anesthetic characteristics and typical ambient concentrations well below the threshold concentration for a biological effect. Nitric oxide is a colorless gas with ambient concentrations generally low enough to have no significant biological effect. Nitrogen dioxide is reddish-brown but is not usually visible at typical ambient concentrations.

2.5.1 Sources

The most significant nitrogen oxide emissions result from the burning of fossil fuels such as coal, oil and gasoline, due to the oxidation of atmospheric nitrogen and nitrogen compounds in the fuel. The primary combustion product is NO , which reacts to form NO_2 .

2.5.2 Effects

At typical concentrations, nitrogen dioxide has significant health effects as a pulmonary irritant, especially upon asthmatics and children. In North Carolina a much greater health concern is the formation of ozone, which is promoted by the presence of NO_2 and other nitrogen oxides.

Some types of vegetation are very sensitive to NO_2 , including oats, alfalfa, tobacco, peas and carrots. Chronic exposure causes chlorosis (yellowing) and acute exposure usually causes irregularly shaped lesions on the leaves.

Nitric oxide and nitrogen dioxide do not directly damage materials. However, NO_2 can react with moisture in the air to produce nitric acid, which corrodes metal surfaces and contributes to acid rain.

High concentrations of NO_2 may reduce visibility. Much of the brownish coloration sometimes observed in polluted air in winter months may be due to NO_2 .

2.6 Lead

Lead is a toxic heavy metal element occurring in the atmosphere as small particles.

2.6.1 Sources

The major source of atmospheric lead used to be the combustion of gasoline containing the additive tetraethyl lead as an antiknock agent. However, the availability of leaded fuel has declined, and the concentration of lead in such fuel has decreased, minimizing gasoline as a source. Significant remaining sources include coal combustion (lead exists

in very small quantities as an impurity in coal) and sandblasting of highway structures and water tanks. Lead also is used in some batteries, paints, insecticides and newspaper inks.

2.6.2 Effects

Lead (Pb) persists and accumulates in the environment and the human body. It may be inhaled, ingested, and eventually absorbed into the bloodstream and distributed to all body tissues. Exposure to low concentrations interferes with blood production and specific enzyme systems. It is believed to cause kidney and nerve cell damage, and severe lead poisoning is known to cause brain damage in children.

3 Standards

Ambient air quality status is determined by measuring pollutant concentrations in outdoor air and comparing the measured concentrations to corresponding standards. The US EPA (Environmental Protection Agency) defines the ambient air as that portion of the atmosphere, external to buildings, to which the general public has access.≡

Ambient air quality standards are classified as primary and secondary. Primary standards are those established to protect public health. Secondary standards are those established to protect the public welfare from adverse pollution effects on soils, water, crops, vegetation, manmade

materials, animals, wildlife, weather, visibility, climate, property, transportation, economy, and personal comfort and well-being. The scientific criteria upon which the standards are based are reviewed periodically by the EPA, which may reestablish or change the standards according to its findings.

A pollutant measurement that is greater than the ambient air quality standard for a specific averaging time is called an exceedance.≡ The national primary, secondary and North Carolina ambient air quality standards are summarized in Table 3.1.

Table 3.1 National and North Carolina Ambient Air Quality Standards

For new or anticipated new standards, References in the Code of Federal Regulations are given. For standards expressed in parts per million, an equivalent mass per unit volume is also shown.

Pollutant/ Ambient Measurement/ (Reference)	Averaging Period	Type of Summary	Primary National (Health Related) Standard	Secondary National (Welfare Related) Standard	North Carolina Standard
TSP 24 hour average	1 year	geometric mean	(1)	(1)	75 µg/m ³
	1 day	2nd maximum	(1)	(1)	150 µg/m ³
PM-2.5 24 hour average (40CFR50, App. N)	1 year	average ² arithmetic mean	15 µg/m ³ (6)	15 µg/m ³ (6)	15 µg/m ³ (6)
	1 day	average ² 98th percentile	65 µg/m ³	65 µg/m ³	65 µg/m ³ (6)
PM-10 24 hour average (40CFR50, App. N)	1 year	average ² arithmetic mean	50 µg/m ³	50 µg/m ³	50 µg/m ³
	1 day	average ² 99th percentile ³	150 µg/m ³	150 µg/m ³	150 µg/m ³
CO 1 hour average	8 hours	2nd maximum	9 ppm (10 mg/m ³)		9 ppm (10 mg/m ³)
	1 hour	2nd maximum	35 ppm (40 mg/m ³)		35 ppm (40 mg/m ³)
O ₃ 1 hour average (40CFR50, App. I)	1 hour	expected ⁴ 2nd maximum	0.12 ppm (6) (235 µg/m ³)	0.12 ppm (6) (235 µg/m ³)	0.12 ppm (235 µg/m ³) (6,7)
	8 hours	average ⁵ arithmetic mean 4th maximum	0.08 ppm (6) (157 µg/m ³)	0.08 ppm (6) (157 µg/m ³)	0.08 ppm (6) (157 µg/m ³)
SO ₂ 1 hour average	1 year	arithmetic mean	0.03 ppm (80 µg/m ³)		0.03 ppm (80 µg/m ³)
	1 day	2nd maximum	0.14 ppm (365 µg/m ³)		0.14 ppm (365 µg/m ³)
	3 hours (non-overlapping)	2nd maximum		0.50 ppm (1,300 µg/m ³)	0.50 ppm (1,300 µg/m ³)
NO ₂ 1 hour average	1 year	arithmetic mean	0.053 ppm (100 µg/m ³)	0.053 ppm (100 µg/m ³)	0.053 ppm (100 µg/m ³)
Pb 24-hour average	1 quarter	arithmetic mean	1.5 µg/m ³	1.5 µg/m ³	1.5 µg/m ³

1. In 1987, National standards for PM-10 replaced those for TSP.

2. Arithmetic mean over the 3 most current years.

3. In July 1997, this percentile-based statistic replaced the 2nd maximum.

4. Determined by adjusting for incomplete days and averaging over the most recent 3 consecutive, complete calendar years.

5. Arithmetic mean value over the most recent 3 consecutive, complete calendar years.

6. On April 1, 2000 North Carolina adopted the EPA PM2.5 and Ozone standards. On May 14, 2000 the US Court of Appeals ruled the new EPA PM2.5 standard vacated and the new 8-hour ozone standard as unenforceable. On appeal to the US Supreme Court the new standard was upheld.

7. On May 27, 2000, the one-hour ozone standard was rescinded by the Environmental Management Commission based on EPA guidance. The one-hour standard is being reinstated by EPA.

4 Ambient Air Quality Monitoring Program

Ambient monitoring and analyses of samples were conducted by the North Carolina Division of Air Quality, three local air pollution control programs and one tribal program.(Appendix A, pp.61-63). The air monitoring data are used to determine whether air quality standards are being met, to assist in enforcement actions, to determine the improvement or decline of air quality, and to determine the extent of allowable industrial expansion, and to provide air pollution information to the public. A list of all monitoring sites active in 2000 is presented in Table 4.1 and shown as a map in Figure 4.1. The locations of sites for individual pollutants are shown in Figures 5.1, 5.4, 5.7, 5.10, 5.16, and 5.19.

Siting of monitors involves several considerations, including size of the area represented, distance from roadways and nearby sources, unrestricted air flow, safety, availability of electricity and security. Each site has a defined monitoring objective, and annual evaluations are conducted to ensure that the objectives are met. The four basic monitoring objectives are to determine:

- # the highest concentration expected in an area;
- # representative concentrations in areas of high population density;
- # the impact of significant sources or source categories on ambient air quality;
- # general background concentration levels.

All monitors have known precision, accuracy, interferences and operational parameters. The monitors B as well as all measurement devices B are carefully calibrated at predetermined frequencies, varying from daily to quarterly. Measurements are traceable to National Institute of Standards and Technology (NIST), when standards are available.

Monitoring and analyses are performed according to a set of standard operating procedures. Field personnel visit manual sampling sites once every six days to replace sample media and check the operation and calibration of monitors. Personnel check continuous monitors at least twice monthly for correct instrument operation.

Quality assurance activities are carried out to determine the quality of the collected ambient data, improve the quality of the data and evaluate how well the monitoring system operates. The goal of quality assurance activities is to produce high quality air pollution data with defined completeness, precision, accuracy, representativeness and comparability.

Microprocessors are used at most sites to collect the data. A computerized telemetry system aids in assembly of the data for submission to the US EPA. This enhances data validity, minimizes travel costs, and allows real-time data to be available by computer polling when needed. Numerous checks are performed to ensure that only valid data are reported.




Figure 4.1 Monitoring Sites Active in 2000

Table 4.1 Ambient Air Monitoring Sites Operated in North Carolina, 2000

SITE COUNTY	ADDRESS	POLLUTANTS	
37-001-0002 ALAMANCE	827 S GRAHAM & HOPE BURLINGTON	PM2.5	
37-003-0003 ALEXANDER	STATE ROAD 1177 TAYLORSVILLE	O3	
37-011-0002 AVERY	7510 BLUE RIDGE PARKWAY	O3	
37-013-0003 BEAUFORT	NC HIGHWAY 306	SO2	
37-021-0003 BUNCOMBE	HEALTH & SOCIAL SERVICES BLDG WOODFIN ST ASHEVILLE	PM10	
37-021-0030 BUNCOMBE	ROUTE 191 SOUTH BREVARD ROAD ASHEVILLE	O3	
37-021-0034 BUNCOMBE	175 BINGHAM ROAD ASHEVILLE	PM2.5	
37-025-0004 CABARRUS	FLOYD STREET KANNAPOLIS	PM10	PM2.5
37-027-0003 CALDWELL	HIGHWAY 321 NORTH LENOIR	O3	
37-029-0099 CAMDEN	COUNTY ROAD 1136 & 1134	O3	
37-033-0001 CASWELL	CHERRY GROVE RECREATION	O3	PM2.5
37-035-0004 CATAWBA	1650 1ST STREET HICKORY	PM10	PM2.5
37-035-0005 CATAWBA	7075 WEST NC HWY 10 HICKORY	PM2.5	
37-037-0004 CHATHAM	ROUTE 4 BOX 62 PITTSBORO	O3	
37-051-0004 CUMBERLAND	F.S. # 5 3296 VILLAGE DRIVE FAYETTEVILLE	PM10	

37-051-0007 CUMBERLAND	CUMBERLAND CO ABC BOARD 1705 OWEN DRIVE FAYETTEVILLE	CO					
37-051-0008 CUMBERLAND	1/4 MILE SR1857/US301/1857	O3					
37-051-0009 CUMBERLAND	4533 RAEFORD ROAD FAYETTEVILLE	PM10	PM2.5				
37-051-1003 CUMBERLAND	3625 GOLFFVIEW ROAD HOPE MILLS	O3					
37-057-0002 DAVIDSON	SOUTH SALISBURY STREET LEXINGTON	PM2.5					
37-057-1002 DAVIDSON	400 SALEM STREET THOMASVILLE	PM10					
37-059-0002 DAVIE	246 MAIN STREET COOLEEMEE	O3					
37-061-0002 DUPLIN	HIGHWAY 50 KENANSVILLE	O3	PM2.5				
37-063-0001 DURHAM	HEALTH DEPT 300 E MAIN ST DURHAM	PM10	PM2.5				
37-063-0011 DURHAM	210 NORTH ROXBORO STREET DURHAM	CO					
37-063-0013 DURHAM	2700 NORTH DUKE STREET DURHAM	O3	HSCO	NOy			
37-065-0003 EDGEcombe	TALBERT PARK at SPRUCE STREET ROCKY MOUNT	PM10	PM2.5				
37-065-0099 EDGEcombe	RT 2, BOX 195 TARBORO	O3					
37-067-0022 FORSYTH	1300 BLK. HATTIE AVENUE WINSTON-SALEM	O3	HSCO	SO2	NO2,y	PM10	PM2.5
37-067-0023 FORSYTH	1401 CORPORATION PARKWAY WINSTON-SALEM	CO	PM10				
37-067-0024 FORSYTH	NORTH FORSYTH HIGH SCHOOL WINSTON-SALEM	PM2.5					
37-067-0025 FORSYTH	100 SW STRATFORD ROAD WINSTON-SALEM	CO					

37-067-0027 FORSYTH	7635 HOLLYBERRY LANE WINSTON-SALEM	O3		
37-067-0028 FORSYTH	6496 BAUX MOUNTAIN ROAD WINSTON-SALEM	O3		
37-067-0029 FORSYTH	1985 GRIFFITH ROAD WINSTON-SALEM	CO		
37-067-1008 FORSYTH	3656 PIEDMONT MEMORIAL DRIVE WINSTON-SALEM	O3	NOy	
37-069-0001 FRANKLIN	431 S. HILLSBOROUGH STREET FRANKLINTON	O3	NOy	
37-071-0016 GASTON	1622 EAST GARRISON BLVD GASTONIA	PM10	PM2.5	
37-077-0001 GRANVILLE	WATER TREATMENT PLANT JOHN UMSTEAD HOSPITAL BUTNER	O3	HSCO	NOy
37-081-0009 GUILFORD	EDGEWORTH & BELLEMEADE ST'S GREENSBORO	PM10	PM2.5	
37-081-0011 GUILFORD	KEELY PARK, KEELY ROAD MC CLEANSVILLE	O3		
37-081-1005 GUILFORD	E. GREEN & S. CENTENNIAL ST HIGH POINT	PM10	PM2.5	
37-081-1011 GULFORD	401 WEST WENDOVER GREENSBORO	CO		
37-085-0001 HARNETT	MUNICIPAL BUILDING DUNN	PM10		
37-087-0002 HAYWOOD	CANTON FIRE DEPARTMENT CANTON	PM10		
37-087-0004 HAYWOOD	2177 ASHEVILLE ROAD WAYNESVILLE	O3		
37-087-0010 HAYWOOD	9 MAIN STREET WAYNESVILLE	PM2.5		
37-087-0035 HAYWOOD	TOWER BLUE RODGE PARKWAY MILE MARKER 410	O3		
37-087-0036 HAYWOOD	GREAT SMOKY MOUNTAIN NATIONAL PARK	O3		
37-089-1006 HENDERSON	CORNER OF ALLEN & WASHINGTON ST'S HENDERSONVILLE	PM10		

37-099-0005 JACKSON	BARNET KNOB FIRE TOWER	O3					
37-099-0006 JACKSON	US ROUTE 19 NORTH CHEROKEE RESERVATION	PM2.5					
37-101-0002 JOHNSTON	3411 JACK ROAD CLAYTON	O3					
37-107-0004 LENIOR	CORNER HWY 70 EAST KINSTON	O3	PM2.5	NOy			
37-109-0004 LINCOLN	RIVERVIEW ROAD LINCOLNTON	O3	NOy	SO2			
37-111-0004 MC DOWELL	BALDWIN AVE MARION	PM10	PM2.5				
37-117-0001 MARTIN	HAYES STREET #2 WELL SITE	O3					
37-119-0001 MECKLENBURG	600 EAST TRADE STREET CHARLOTTE	TSP	PM10				
37-119-0003 MECKLENBURG	FIRE STATION # 11, 620 MORETZ ST CHARLOTTE	PM10					
37-119-0010 MECKLENBURG	FIRE STATION # 10, 2136 FREMOUNT RD CHARLOTTE	TSP	PM10	PM2.5			
37-119-0035 MECKLENBURG	1330 SPRING STREET GRANVILLE NEIGHBORHOOD CENTER CHARLOTTE	CO					
37-119-0038 MECKLENBURG	301 TRYON STREET CHARLOTTE	CO					
37-119-0040 MECKLENBURG	6623 PARK SOUTH DRIVE CHARLOTTE	PM2.5					
37-119-0041 MECKLENBURG	2210 EASTWAY DRIVE CHARLOTTE	O3	HSCO	CO	SO2	NO2,y	PM2.5
37-119-0042 MECKLENBURG	1935 EMERYWOOD DRIVE CHARLOTTE	PM2.5					
37-119-1001 MECKLENBURG	FILTER PLANT DAVIDSON	PM10					
37-119-1005 MECKLENBURG	400 WESTINGHOUSE BLVD. CHARLOTTE	O3	PM10				

37-119-1009 MECKLENBURG	29N @ MECKLENBURG CAB CO. CHARLOTTE	O3	HSCO	NOy
37-121-0001 MITCHELL	CITY HALL, SUMMIT STREET SPRUCE PINE	PM10	PM2.5	
37-123-0001 MONTGOMERY	112 PERRY DRIVE	PM2.5		
37-129-0002 NEW HANOVER	6028 HOLLY SHELTER ROAD	O3		
37-129-0006 NEW HANOVER	HIGHWAY 421 NORTH WILMINGTON	SO2		
37-129-0008 NEW HANOVER	CORNER of OLEANDER & COLLEGE WILMINGTON	CO		
37-129-0009 NEW HANOVER	2710 MARKET STREET WILMINGTON	PM10	PM2.5	
37-131-0002 NORTHAMPTON	ROUTE 46 GASTON	O3		
37-133-0005 ONSLow	617 HENDERSON DRIVE JACKSONVILLE	PM10	PM2.5	
37-135-0006 ORANGE	147 EAST FRANKLIN STREET CHAPEL HILL	CO		
37-135-0007 ORANGE	MASON FARM ROAD CHAPEL HILL	PM2.5		
37-139-0002 PASQUOTANK	600 WESTOVER STREET ELIZABETH CITY	PM10	PM2.5	
37-145-0003 PERSON	STATE ROAD 1102 & NC 49	O3		
37-147-0005 PITT	851 HOWELL STREET GREENVILLE	PM10	PM2.5	
37-147-0099 PITT	US 264 NEAR WATER TOWER FARMVILLE	O3	SO2	
37-155-0004 ROBESON	1110 MIMOSA STREET LUMBERTON	PM2.5		
37-155-0004 ROBESON	1170 LINKHAW ROAD LUMBERTON	PM2.5		

37-157-0099 ROCKINGHAM	6371 NC 65 @ BETHANY SCHOOL BETHANY	O3			
37-159-0021 ROWAN	WEST STREET & GOLD HILL AVENUE ROCKWELL	O3	HSCO	NOy	
37-159-0022 ROWAN	925 NORTH ENOCHVILLE AVENUE	O3	HSCO	NOy	
37-163-0003 SAMPSON	BREWER ROAD FAISON	NOy			
37-163-0004 SAMPSON	BREWER ROAD FAISON	NOy			
37-163-0005 SAMPSON	HORTICULTURAL CROPS FAISON	NOy			
37-173-0002 SWAIN	CENTER STREET, PARKS 7 RECREATION FACILITY	O3	PM10	PM2.5	
37-179-0002 UNION	701 CHARLES STREET MONROE	O3			
37-183-0003 WAKE	FIRE STATION #9 SIX FORKS ROAD RALEIGH	PM10			
37-183-0011 WAKE	420 PERSON STREET RALEIGH	CO			
37-183-0014 WAKE	EAST MILLBROOK JR HI 3801 SPRING FOREST ROAD RALEIGH	O3	PM10	PM2.5	
37-183-0015 WAKE	808 NORTH STATE STREET RALEIGH	O3	HSCO	PM2.5	NOy
37-183-0016 WAKE	201 NORTH BROAD STREET FUQUAY-VARINA	O3			
37-183-0017 WAKE	5033 TV TOWER ROAD GARNER	O3			
37-191-0005 WAYNE	DILLARD MIDDLE SCHOOL GOLDSBORO	PM10			
37-199-0003 YANCEY	BLUE RIDGE PARKWAY	O3			

Sites Operated in 2000	100
---------------------------	-----

5 Pollutant Monitoring Results

Air quality in a given area is affected by many factors, including meteorological conditions, the location of pollutant sources, and the amount of pollutants emitted from them.

The speed and direction of air movement determine whether pollutant emissions cause exceedances of the ambient air quality standards and where those exceedances will occur. Atmospheric stability, precipitation, solar radiation and temperature also affect pollutant concentrations.

Geographic factors that affect concentrations include variables such as whether an area is urban or rural, and whether the area has mountains, valleys or plains.

Important economic factors affecting air quality include concentration of industries, conditions of the economy, and the day of the week.

Air quality also may be influenced by exceptional events in the short term. Exceptional events may be either natural (e.g., forest fire) or manmade (e.g., construction or demolition). Unusual data that can be attributed to an exceptional event are considered biased and may be omitted

from data summaries when they are not representative of normal conditions. In the tabular listings in this report, data affected by exceptional events are excluded, and are omitted from summaries in charts. However they are addressed in the text of the report. A list of typical exceptional events is given in Appendix B.

Data for the 2000 ambient air quality report were collected at 173 air pollutant monitors operated by state and local agencies in North Carolina (listed in Appendix A, pp. 61-63). To minimize operating expenses, some sulfur dioxide monitors are operated only every third year. Sixteen of the 173 monitors used for this report operated most recently in 1998 or 1999.

5.1 Total Suspended Particulates

Total Suspended Particulate matter (TSP) is collected on filters using a high volume sampler (an EPA Reference Method). The sampler motor is set and calibrated to an air flow rate of 40 ± 4 cubic feet per minute. Gravimetric analysis is performed by comparing the exposed filter weight to the unexposed filter weight. Weights are

measured to the nearest 0.1 milligram. The difference between the exposed and unexposed weights is the amount of particulate collected from a known volume of air.

In 2000, two sites in Mecklenburg County were used to monitor TSP and 92 samples were collected. A detailed summary of the data from each site is given in Table 5.1.

No sample exceeded the N.C. TSP ambient air quality standards in 2000. The highest 24-hour average was 116, which was 77 percent of the standard. This value occurred at the Fire Station # 10 site in Charlotte. Attainment status is based on the second highest 24-hour concentration and on the

geometric mean of all the 24-hour concentrations at a given site.

The largest geometric mean TSP average was $45 \mu\text{g}/\text{m}^3$, which is 59 percent of the level of the air quality standard. This value occurred at the East Trade Street site in Charlotte.

During early November 2000, especially November 2, 3, 4, and 8 there were several sites on which particulate matter samplers were affected by smoke from forest fires in North Carolina and neighboring states. A reading of $152 \mu\text{g}/\text{m}^3$, which is above the standard of $150 \mu\text{g}/\text{m}^3$, was recorded at the East Trade Street site in Charlotte. The geometric mean increased to $49 \mu\text{g}/\text{m}^3$ as well.

Table 5.1 Total Suspended Particulates in Micrograms Per Cubic Meter for 2000

SITE NUMBER COUNTY	ADDRESS	NUM OBS	24- 1ST	HOUR 2ND	MAX 3RD	4TH	ARITH MEAN	GEOM MEAN	GEOM SD
37-119-0001 MECKLENBURG	600 EAST TRADE STREET CHARLOTTE	40	90	82	75	70	47	45	1.4
37-119-0010 MECKLENBURG	FIRE STA #10 2136 REMOUNT ROAD CHARLOTTE	52	116	75	73	71	44	41	1.5
Total Samples		92							
Total Sites Sampled		2							

5.2 PM₁₀

State and local program agencies in North Carolina use high volume samplers and size selective inlets to collect PM₁₀ samples. A gravimetric analysis procedure (EPA Reference Method) is used to analyze the samples.

In 2000, 32 sites were used to monitor PM₁₀ and 2,628 samples were collected. A map of the PM₁₀ sampling sites is shown in Figure 5.1, and a detailed summary of the data from each site is given in Table 5.2.

There were no exceedances of the PM₁₀ ambient air quality standards in 2000. The highest 24-hour maximum concentration was 98 µg/m³, or about 65 percent of the standard (150 µg/m³). This value occurred at

the Westinghouse Blvd site in Charlotte. The highest annual arithmetic mean was 31 µg/m³, which is 62 percent of the standard (50 µg/m³). This annual average occurred at the same site in Mecklenburg County. The previously described fires did cause a value of 113 µg/m³ to occur in Canton, Haywood County.

The second highest 24-hour concentrations are shown by county in Figure 5.2 and the annual arithmetic means are shown in Figure 5.3. (In counties with more than one PM₁₀ monitoring site, the concentration reported in Figure 5.2 is the county-wide second maximum 24-hour concentration, and the mean reported in Figure 5.3 is the maximum arithmetic mean for the county.)




Figure 5.1 Location of PM₁₀ Monitoring Sites

Table 5.2 PM10 in Micrograms Per Cubic Meter for 2000

SITE NUMBER COUNTY	ADDRESS	NUM OBS	24-HOUR				ARITH MEAN
			1ST	2ND	3RD	4TH	
37-021-0003 BUNCOMBE	HEALTH & SOCIAL SERVICES BLDG WOODFIN ST ASHEVILLE	59	41	38	38	36	18
37-025-0004 CABARRUS	FLOYD STREET KANNAPOLIS	56	47	40	38	37	19
37-035-0004 CATAWBA	1650 1ST. ST. HICKORY	55	47	42	40	37	22
37-051-0004 CUMBERLAND	F.S. # 5 3296 VILLAGE DR. FAYETTEVILLE	41	78	51	42	39	28
37-051-0009 CUMBERLAND	4533 RAEFORD ROAD FAYETTEVILLE	59	63	52	41	38	23
37-057-1002 DAVIDSON	400 SALEM STREET THOMASVILLE	60	43	41	38	37	21
37-063-0001 DURHAM	HEALTH DEPT 300 E MAIN ST DURHAM	58	44	43	43	43	21
38-065-0003 EDGECOMBE	TALBERT PARK at SPRUCE ST ROCKY MOUNT	56	43	41	33	33	20
37-067-0022 FORSYTH	1300 BLK. HATTIE AVE WINSTON-SALEM	364	50	47	47	47	21
37-067-0023 FORSYTH	1401 CORPORATION PARKWAY WINSTON-SALEM	296	54	51	50	49	22
37-071-0016 GASTON	1622 E. GARRISON BLVD GASTONIA	56	40	37	36	36	21
37-081-0009 GUILFORD	EDGEWORTH & BELLEMEADE GREENSBORO	60	54	44	42	40	23
37-081-1005 GUILFORD	E GREEN & S CENTENNIAL ST HIGH POINT	56	45	44	44	44	24
37-085-0001 HARNETT	MUNICIPAL BUILDING DUNN	59	52	52	48	44	28
37-087-0002 HAYWOOD	ROOF, CANTON FIRE DEPT. CANTON	60	51	47	46	44	26
37-089-1006 HENDERSON	CORNER OF ALLEN & WASHINGTON STS HENDERSONVILLE	54	50	44	43	40	23
37-111-0004 MC DOWELL	BALDWIN AVE & EAST M STREET MARION	58	46	45	39	37	22
37-119-0001 MECKLENBURG	600 EAST TRADE STREET CHARLOTTE	31	48	45	43	42	27
37-119-0003 MECKLENBURG	FIRE STA #11 620 MORETZ STREET	58	97	62	56	51	31

SITE NUMBER COUNTY	ADDRESS	NUM OBS	24-HOUR				ARITH MEAN
			1ST	2ND	3RD	4TH	
37-119-0010 MECKLENBURG	CHARLOTTE FIRE STA #10 2136 REMOUNT ROAD CHARLOTTE	56	79	48	48	46	29
37-119-1001 MECKLENBURG	FILTER PLANT DAVIDSON	53	63	54	50	46	26
37-119-1005 MECKLENBURG	400 WESTINGHOUSE BLVD. CHARLOTTE	41	98	62	61	59	31
37-121-0001 MITCHELL	CITY HALL SUMMIT ST SPRUCE PINE	60	88	50	48	45	27
37-129-0009 NEW HANOVER	2710 MARKET STREET WILMINGTON	60	45	36	34	33	17
37-133-0005 ONslow	617 HENDERSON DRIVE JACKSONVILLE	60	33	32	31	28	17
37-139-0002 PASQUOTANK	600 WESTOVER STREET ELIZABETH CITY	57	38	34	32	30	17
37-147-0003 PITT	851 HOWELL STREET GREENVILLE	58	39	36	36	33	19
37-173-0002 SWAIN	CENTER ST/PARKS 7 REC FACILITY	58	46	33	32	32	19
37-183-0003 WAKE	FIRE STATION #9 SIX FORKS RD NORTH HILLS RALEIGH	60	51	42	39	39	23
37-183-0014 WAKE	E MILLBROOK JR HI RALEIGH	356	61	51	50	50	22
37-183-0014 WAKE	E MILLBROOK JR HI RALEIGH	55	45	40	38	35	22
37-191-0004 WAYNE	DILLARD MIDDLE SCHOOL	58	41	40	39	33	21
Total Samples		2628					
Total Sites Sampled		32					




Figure 5.2 PM₁₀: Second Highest 24-Hour Averages, 2000




Figure 5.3 PM₁₀: Maximum Annual Arithmetic Means, 2000

5.3 $PM_{2.5}$

In 2000, 38 sites were used to monitor $PM_{2.5}$ and 5,188 samples were collected. A map of the $PM_{2.5}$ sampling sites is shown in Figure 5.4 and a detailed summary of the data from each site is given in Table 5.3.

There was one exceedance of the $PM_{2.5}$ 24-hour ambient air quality standards in 2000. The highest 24-hour maximum concentration was $67.4 \mu\text{g}/\text{m}^3$, or about 104 percent of the standard ($65 \mu\text{g}/\text{m}^3$). This value occurred at the Raeford Road site in Fayetteville. However the forest fires caused 5 values above the standard to be flagged. These values of 84.5, 84.3, 84.2, 82.7 and $69.4 \mu\text{g}/\text{m}^3$ occurred Charlotte, Marion,

Charlotte, Waynesville and Winston-Salem respectively.

The highest annual arithmetic mean was $17.8 \mu\text{g}/\text{m}^3$, which is 119 percent of the standard ($15 \mu\text{g}/\text{m}^3$). This annual average occurred at the South Salisbury Street site in Lexington, Davidson County. However 27 of the 38 monitors also exceeded the annual arithmetic mean standard.

The second highest 24-hour concentrations are shown by county in Figure 5.6 and the annual arithmetic means are shown in Figure 5.7. (In counties with more than one $PM_{2.5}$ monitoring site, the concentration reported in Figure 5.6 is the county-wide second maximum 24-hour concentration, and the mean reported in Figure 5.7 is the maximum arithmetic mean for the county.)




Figure 5.4 Location of $PM_{2.5}$ Monitoring Sites

Table 5.3 PM_{2.5} in Micrograms Per Cubic Meter for 2000

SITE NUMBER COUNTY	ADDRESS	NUM OBS	24-HOUR MAXIMA				ARITH MEAN
			1ST	2ND	3RD	4TH	
37-021-0003 ALAMANCE	827 SOUTH GRAHAM & HOPE BURLINGTON	102	34.5	33.8	33.3	36.6	15.3
37-021-0034 BUNCOMBE	175 BINGHAM ROAD ASHEVILLE	78	36.8	34.1	32.9	32.0	15.8
37-025-0004 CABARRUS	FLOYD STREET KANNAPOLIS	110	42.5	33.0	32.5	31.3	16.6
37-033-0001 CASWELL	CHERRY GROVE RECREATION	115	45.8	36.2	35.0	32.9	14.9
37-035-0004 CATAWBA	1650 1ST. ST. HICKORY	112	38.0	36.7	34.2	33.5	17.4
37-035-0005 CATAWBA	7075 WEST HWY 10 HICKORY	68	35.8	34.1	31.7	31.5	16.7
37-037-0004 CHATHAM	RT4 BOX62 PITTSBORO	115	32.2	27.9	26.8	26.6	13.3
37-051-0009 CUMBERLAND	4533 RAEFORD ROAD FAYETTEVILLE	109	67.4	50.2	33.0	31.4	15.9
37-057-0002 DAVIDSON	SOUTH SALISBURY STREET LEXINGTON	103	46.8	37.7	34.9	34.1	17.8
37-061-0001 DUPLIN	HWY 50 KENANANSVILLE	117	33.1	31.4	24.7	24.7	13.1
37-063-0001 DURHAM	HEALTH DEPT 300 E MAIN ST DURHAM	351	43.0	39.6	37.4	30.8	14.7
38-065-0003 EDGEcombe	TALBERT PARK at SPRUCE ST ROCKY MOUNT	112	34.9	31.7	30.8	28.0	14.7
37-067-0022 FORSYTH	1300 BLOCK, HATTIE AVENUE WINSTON-SALEM	350	44.2	38.5	37.5	37.4	16.5
37-067-0024 FORSYTH	NORTH FORSYTH HIGH SCHOOL WINSTON-SALEM	105	43.2	35.3	35.0	34.07	15.9
37-071-0016 GASTON	1622 EAST GARRISON BLVD GASTONIA	119	36.8	31.0	29.8	29.8	15.9
37-081-0009 GUILFORD	EDGEWORTH & BELLEMEADE GREENSBORO	318	52.3	37.2	36.2	35.0	16.7
37-081-1005 GUILFORD	E GREEN & S CENTENNIAL ST HIGH POINT	104	39.2	34.4	33.5	33.0	16.1
37-087-0010 HAYWOOD	9 MAIN STREET WAYNESVILLE	110	33.8	33.0	31.8	28.9	14.7
37-099-0006 JACKSON	US RT 19 NORTH CHEROKEE RES	80	33.7	29.8	29.6	28.6	14.0

SITE NUMBER COUNTY	ADDRESS	NUM OBS	24-HOUR				ARITH MEAN
			1ST	2ND	3RD	4TH	
37-107-0004 LENOIR	CORNER HWY 70 EAST KINSTON	115	34.2	32.4	24.3	22.7	12.7
37-111-0004 MC DOWELL	BALDWIN AVE MARION	114	39.1	38.5	34.7	34.3	16.4
37-119-0010 MECKLENBURG	FIRE STA #10 2136 REMOUNT ROAD CHARLOTTE	339	46.9	38.5	3805	36.9	17.0
37-119-1040 MECKLENBURG	6623 PARK SOUTH DRIVE CHARLOTTE	81	45.2	32.8	31.2	28.9	16.0
37-119-1041 MECKLENBURG	1120 EASTWAY DRIVE CHARLOTTE	341	43.5	36.0	35.4	35.6	16.3
37-119-1042 MECKLENBURG	1935 EMERYWOOD DRIVE CHARLOTTE	33	37.8	29.4	28.7	27.8	17.6
37-121-0001 MITCHELL	CITY HALL SUMMIT ST SPRUCE PINE	110	42.6	37.2	35.4	33.2	15.1
37-123-0001 MONTGOMERY	112 PERRY DRIVE	101	60.9	24.1	28.3	26.4	14.2
37-129-0009 NEW HANOVER	2710 MARKET STREET WILMINGTON	111	31.7	30.2	28.0	24.9	12.4
37-133-0005 ONSLOW	617 HENDERSON DRIVE JACKSONVILLE	117	34.3	30.7	27.7	25.0	12.3
37-135-0007 ORANGE	MASON FARM ROAD CHAPEL HILL	117	30.2	29.7	29.7	28.6	14.4
37-139-0002 PASQUOTANK	600 WESTOVER STREET ELIZABETH CITY	104	37.3	32.8	28.3	26.6	12.3
37-147-0005 PITT	851 HOWELL STREET GREENVILLE	115	41.0	33.3	30.5	28.5	13.9
37-155-0004 ROBESON	1110 MIMOSA STREET LUMBERTON	29	21.4	20.9	20.6	20.1	13.9
37-155-0005 ROBESON	1170 LINKHAM ROAD LUMBERTON	12	26.3	21.9	20.1	18.0	15.3
37-173-0002 SWAIN	CENTER ST/PARKS 7 REC FACILITY	107	37.7	33.6	31.9	31.3	14.2
37-183-0014 WAKE	EAST MILLBROOK JR HI RALEIGH	345	52.8	40.2	37.7	35.3	15.8
37-183-0015 WAKE	808 NORTH STATE STREET RALEIGH	105	40.2	32.0	29.6	28.0	14.4
37-191-0005 WAYNE	DILLARD MIDDLE SCHOOL GOLDSBORO	113	39.7	38.2	34.4	30.5	15.9
Total Samples		5188					
Total Sites Sampled		38					




Figure 5.5 PM_{2.5}: Second Highest 24-Hour Averages, 2000




Figure 5.6 PM_{2.5}: Maximum Annual Arithmetic Means, 2000

5.4 Carbon Monoxide

Carbon monoxide (CO) data were collected for two purposes in 2000: to determine attainment status of the ambient air quality standard and to gather data on CO as an ozone precursor. The carbon monoxide associated with ozone formation consists of very low concentrations (not greater than 2 ppm) collected at special sites considered optimal for input to a large photochemical grid model. This report will not further discuss the role of CO as an ozone precursor, but these data and more information are available on request from the Division of Air Quality (see the Preface for a mailing address).

To assess CO attainment status, the Division of Air Quality collected data from monitors in Fayetteville, Chapel Hill, Wilmington, Durham, Greensboro and Raleigh, and local program agencies collected data from three monitors in Winston-Salem and Charlotte using EPA Reference or equivalent methods to measure the concentrations.

In 2000, 12 sites were used to monitor CO and 61,959 valid hourly averages were collected. To keep operating costs minimal, some sites are operated only in the colder months. A map of the CO sampling sites is shown in Figure 5.7, and a detailed summary of the data from each site is presented in Table 5.4.

There were no exceedances of the CO ambient air quality standards in 1998. The highest 1-hour average was 14.1 parts per million (ppm), or about 40 percent of the standard (35 ppm). This value occurred at the Tryon Street site in Charlotte. The highest 8-hour average was 5.4 ppm, at the

same site, which is 60 percent of the standard.

The second highest 1-hour concentrations in each county are shown in Figure 5.8 and the second highest 8-hour concentrations are shown in Figure 5.9.

Historical data have demonstrated that high concentrations of CO occur more frequently in autumn and winter than during the warmer months of the year. There are three main reasons for this seasonal variation: (1) North Carolina experiences more atmospheric inversions in colder months, trapping air pollutants at low heights; (2) motor vehicles emit more CO due to inefficient combustion during cold starts and warm up; and (3) during colder temperatures, more fuel is burned for comfort heating.

All areas monitored are attaining the ambient air quality standards for carbon monoxide. Several factors have reduced CO concentrations, with the most significant being that older vehicles are gradually being replaced with newer, more efficient vehicles. The motor vehicle Inspection and Maintenance program (in effect in Mecklenburg, Wake, Durham, Forsyth, Guilford, Gaston, Cabarrus, Orange and Union counties) is an intentional control strategy that helps assure cleaner-running cars. Other factors include increased news media interest and public awareness, and the reporting of the Air Quality Index (see Chapter 6 of this report). As a result of greater public awareness, more cars are kept in better running condition, thus operating more cleanly. Traffic flow improvements such as new roads and better coordinated traffic signals also help reduce CO.




Figure 5.7 Location of Carbon Monoxide Monitoring Sites

Table 5.4 Carbon Monoxide in Parts Per Million for 2000

SITE NUMBER COUNTY	ADDRESS	NUM OBS	ONE-HOUR MAXIMA		EIGHT-HOUR MAXIMA	
			1ST	2ND	1ST	2ND
37-051-0007 CUMBERLAND	ABC BOARD, 1705 OWEN DR FAYETTEVILLE	4,367	7.3	6.3	5.0	3.9
37-063-0011 DURHAM	201 NORTH ROXBORO ST DURHAM	2,873	6.2	6.1	5.0	4.5
37-067-0023 FORSYTH	1401 CORPORATION PKY WINSTON-SALEM	8,706	4.8	4.7	4.0	3.3
37-067-0025 FORSYTH	100 SW STRATFORD RD WINSTON-SALEM	8,704	2.8	2.8	2.1	1.9
37-067-0029 FORSYTH	1985 GRIFFITH ROAD WINSTON-SALEM	937	5.1	5.1	3.6	3.3
37-081-1001 GILFORD	401 WEST WENDOVER GREENSBORO	4,375	6.0	4.6	4.1	3.3
37-119-0035 MECKLENBURG	1330 SPRING ST GRANVILLE NEIGHBORHOOD CHARLOTTE	3,210	5.0	5.0	4.4	3.5
37-119-0038 MECKLENBURG	301 N TRYON ST CHARLOTTE	8,659	14.1	10.8	4.9	4.7
37-119-0041 MECKLENBURG	1120 EASTWAY DRIVE CHARLOTTE	8,630	5.2	4.8	3.8	3.8
37-129-0008 NEW HANOVER	OLEANDER & COLLEGE WILMINGTON	4,342	7.4	6.3	5.4	3.8
37-135-0006 ORANGE	147 EAST FRANKLIN STREET CHAPEL HILL	2,156	5.8	5.4	4.1	3.8
37-183-0011 WAKE	420 S PERSON ST RALEIGH	5,000	6.8	6.8	5.4	5.2
Total Samples		61,959				
Total Sites Sampled		12				




Figure 5.8 Carbon Monoxide: Second Highest 1-Hour Average, 2000




Figure 5.9 Carbon Monoxide: Second Highest Non-overlapping 8-Hour Average, 2000

5.5 Ozone

Ozone (O₃) concentrations are measured using EPA reference or equivalent continuous monitors. Ozone is a seasonal pollutant formed in the atmosphere as a result of many chemical reactions that occur in sunlight, mainly during the warmer months. Thus, most ozone monitors only operate from April through October.

The state and local program agencies operated 45 monitoring sites in 2000 during the ozone season, April through October. A map of the O₃ sampling sites is presented in Figure 5.10, and a detailed summary of the one-hour data from each site is given in Table 5.5, and the 8-hour data in Table 5.6. These 45 monitoring sites provided 226,922 hourly samples.

There were 8 exceedances of the 1-hour ozone standard in North Carolina in 2000. Durham, Rowan and Wake Counties had one exceedance and Mecklenburg County had the rest.

The one-hour standard is exceeded when one valid one-hour average exceeds 0.124 ppm at a site and the expected number of exceedances is greater than 1. (To exceed the standard, the largest average must be larger than 0.12 ppm when *rounded* to two significant digits. The expected number of exceedances is determined from a 3-year average of exceedance day counts for an area. Moreover, when any ozone sampling day does not have a valid maximum ozone measurement for any reason, the missing day can be counted as an *estimated* exceedance day under certain circumstances [40 CFR 50 App. J, US EPA 1993, p. 767-768]. Table 5.4 gives both the actually

measured and the estimated number of exceedance days at each site.)

Mecklenburg County and the Triad Counties were redesignated as attainment/maintenance areas on July 5, 1995, and November 8, 1993, respectively. Generally, an area is in violation if it exceeds the standard at a monitor four or more times in any three year period. Three sites, one each in Mecklenburg, Rowan and Wake Counties had nine, five and six exceedances in the three-year period 1998-2000. However, since the one-hour standard was rescinded by EPA, and by the EMC on May 27, 2000 the exceedances did not cause the counties to be designated nonattainment. New nonattainment designations based on either the 8-hour standard or the 1-hour standard are likely to be declared after the 2001 sampling seasons are completed and certified. Attainment status as of 2000 is discussed in Appendix D.

The 8-hour standard was exceeded a total of 239 times at the 45 sites that monitored for O₃. Of the 33 counties that monitor for ozone, only four failed to register at least one exceedance. Five counties had 10 or more exceedances at the worst site with Davie County leading the way with 17. These 239 exceedances were distributed over 35 days during the ozone season where at least one site within the state recorded values greater than 0.085 ppm.

The second highest 1-hour concentrations in each county are shown in Figure 5.11 for areas with one or more monitors active in 2000. Figure 5.12 shows the second-highest 8-hour concentrations of O₃ for sites operating in 2000. The level of the 1-hour standard, 0.125 ppm, was exceeded six times in total, on three calendar days (two each on June 1, 2, and 12).

Figure 5.13 shows the distribution of the fourth maximum 8-hour values from 1991 to 2000 as box-and-whisker plots. (See Appendix C for an explanation of this type of chart.) The smoothed regression line of the values is below the standard from 1991 to when it crosses the standard in 1995, rising to a maximum in 1999 and falling off, but still above the standard by 2000. Figure 5.14 shows the number of exceedances of

the 8-hour ozone standard of 0.085 ppm from 1993 to 2000. Figure 5.15 partitions the exceedances by air quality regions. The first column is the Charlotte-Gastonia Metropolitan Statistical Area. The second column is the Greensboro-Winston-Salem-High Point Metropolitan Statistical Area. The third column is the Raleigh-Durham Metropolitan Statistical Area, and the final column combines the rest of the state.




Figure 5.10 Location of Ozone Monitoring Sites

Table 5.5 One-Hour Ozone in Parts Per Million for 2000

SITE NUMBER	ADDRESS	NUM	DAILY 1-HR MAXIMA				NO. VALUES > 0.125 MEAS	EST
			COUNTY	OBS	1ST	2nd		
37-003-0003 ALEXANDER	STATE ROAD 1177 TAYLORSVILLE	5,064	0.117	0.108	0.104	0.098	0	0.0
37-011-0003 AVERY	7510 BLUE RIDGE	4,872	0.090	0.089	0.087	0.086	0	0.0
37-021-0030 BUNCOMBE	ROUT 191 SOUTH BREVARD RD ASHEVILLE	5,112	0.108	0.107	0.106	0.102	0	0.0
37-027-0003 CALDWELL	HWY 321 NORTH LENOIR	4,848	0.099	0.099	0.096	0.095	0	0.0
37-029-0099 CAMDEN	COUNTY ROAD 1136 & 1134	5,040	0.091	0.083	0.079	0.078	0	0.0
37-033-0001 CASWELL	CHERRY GROVE RECREATION	5,112	0.122	0.121	0.112	0.105	0	0.0
37-037-0004 CHATHAM	RT 4 BOX 64 PITTSBORO NC 27312	4,968	0.100	0.099	0.091	0.090	0	0.0
37-051-0008 CUMBERLAND	1/4MI SR1857/US301/1857	5,112	0.104	0.102	0.101	0.100	0	0.0
37-051-1003 CUMBERLAND	3625 GOLFFVIEW ROAD HOPE MILLS	5,064	0.106	0.106	0.101	0.097	0	0.0
37-059-0002 DAVIE	246 MAIN STREET COOLEEMEE	5,112	0.114	0.112	0.111	0.108	0	0.0
37-061-0002 DUPLIN	HWY 50 KENANSVILLE	5,136	0.102	0.095	0.092	0.091	0	0.0
37-063-0013 DURHAM	2700 NORTH DUKE STREET DURHAM	5,112	0.133	0.116	0.112	0.105	1	1.0
37-065-0099 EDGEcombe	RT 2, BOX 195 TARBORO	5,136	0.107	0.106	0.101	0.100	0	0.0
37-067-0022 FORSYTH	1300 BLK. HATTIE AVENUE WINSTON-SALEM	5,136	0.109	0.104	0.099	0.099	0	0.0
37-067-0027 FORSYTH	7635 HOLLYBERRY LANE WINSTON-SALEM	5,112	0.101	0.096	0.095	0.094	0	0.0
37-067-0028 FORSYTH	6496 BAUX MOUNTAIN RD WINSTON-SALEM	5,136	0.112	0.107	0.106	0.096	0	0.0
37-067-1008 FORSYTH	3656 PIEDMONT MEMORIAL DRIVE WINSTON-SALEM	5,112	0.110	0.105	0.103	0.097	0	0.0
37-069-0001 FRANKLIN	431 S. HILLBOROUGH ST FRANKLINTON	5,136	0.113	0.110	0.101	0.099	0	0.0
37-077-0001 GRANVILLE	WATER TREATMENT PLANT JOHN UMSTEAD HOSPITAL BUTNER	5,112	0.113	0.106	0.105	0.102	0	0.0

37-081-0011 GUILFORD	KEELY PARK, KEELY RD, MCCLEANSVILLE	5,122	0.121	0.116	0.109	0.107	0	0.0
37-087-0004 HAYWOOD	2177 SCHEVILLS ROAD WAYNESVILLE	5,136	0.101	0.099	0.094	0.093	0	0.0
37-087-0035 HAYWOOD	TOWER BLUE RIDGE PARKWAY MILE MARKER 410	4,368	0.098	0.093	0.092	0.090	0	0.0
37-087-0036 HAYWOOD	GREAT SMOKY MOUNTAIN NATIONAL PARK	5,136	0.104	0.102	0.092	0.092	0	0.0
37-099-0005 JACKSON	BARNET KNOB FIRE TOWER	4,898	0.105	0.104	0.094	0.092	0	0.0
37-101-0002 JOHNSTON	3411 JACK ROAD CLAYTON	5,122	0.117	0.116	0.097	0.096	0	0.0
37-107-0004 LENIOR	CORNER HWY EAST KINSTON	5,088	0.102	0.097	0.096	0.096	0	0.0
37-109-0004 LINCOLN	RIVERVIEW ROAD LINCOLNTON	5,088	0.122	0.109	0.107	0.103	0	0.0
37-117-0001 MARTIN	HAYES STREET (#2WELL SITE)	5,064	0.096	0.095	0.091	0.091	0	0.0
37-119-0041 MECKLENBURG	1120 EASTWAY DRIVE CHARLOTTE	5,088	0.152	0.130	0.127	0.117	3	3.0
37-119-1005 MECKLENBURG	400 WESTINGHOUSE BLVD. CHARLOTTE	5,136	0.119	0.104	0.104	0.103	0	0.0
37-119-1009 MECKLENBURG	29 N@ MECKLENBURG CAB CO CHARLOTTE	5,088	0.144	0.141	0.121	0.119	2	2.0
37-129-0002 NEW HANOVER	6028 HOLLY SHELTER RD	5,136	0.110	0.098	0.090	0.089	0	0.0
37-131-0002 NORTHAMPTON	ROUTE 46 GASTON	5,112	0.099	0.097	0.096	0.088	0	0.0
37-145-0099 PERSON	SR 1102 & NC 49	5,088	0.137	0.117	0.109	0.109	1	1.0
37-147-0099 PITT	US 264 NEAR WATTER TOWER FARMVILLE	5,136	0.100	0.095	0.095	0.091	0	0.0
37-157-0099 ROCKINGHAM	6371 NC 65 @ BETHANY SCHOOL	5,112	0.105	0.100	0.098	0.092	0	0.0
37-159-0021 ROWAN	WEST ST & GOLD HILL AVENUE ROCKWELL	5,136	0.125	0.122	0.114	0.113	1	1.0
37-159-0022 ROWAN	925 N ENOCHVILLE AVE	5,016	0.123	0.122	0.113	0.113	0	0.0
37-173-0002 SWAIN	CENTER STREET PARKS 7 REC FACILITY	5,064	0.086	0.082	0.082	0.080	0	0.0
37-179-0003 UNION	701 CHARLES STREET MONROE	4,776	0.100	0.095	0.094	0.093	0	0.0
37-183-0014 WAKE	E. MILLBROOK JR HI 3801 SPRING FOREST ROAD RALEIGH	5,136	0.122	0.115	0.114	0.104	0	0.0

37-183-0015 WAKE	808 NORTH STATE STREET RALEIGH	5,136	0.115	0.113	0.107	0.100	0	0.0
37-183-0016 WAKE	201 NORTH BROAD STREET FUQUAY-VARINA	5,112	0.124	0.101	0.101	0.096	0	0.0
37-183-0017 WAKE	5033 TV TOWER ROAD GARNER	5,112	0.135	0.117	0.110	0.098	1	1.0
37-199-0003 YANCY	BLUE RIDGE PARKWAY	4,344	0.111	0.107	0.102	0.102	0	0.0

Total Samples	226,922						8	8.0
Total Sites Sampled	45							

Table 5.6 Eight-Hour Ozone in Parts Per Million for 2000

SITE NUMBER	ADDRESS	NUM	VALID DAILY 8-HOUR MAXIMUM						NO. VALUES >.085 MEAS	
			COUNTY	OBS	1ST	2nd	3rd	4th		5TH
37-003-0003 ALEXANDER	STATE ROAD 1177 TAYLORSVILLE	213		0.099	0.098	0.093	0.091	0.088	0.086	9
37-001-0001 AVERY	7500 BLUE RIDGE	208		0.082	0.080	0.079	0.078	0.078	0.077	0
37-021-0030 BUNCOMBE	ROUT 191 SOUTH BREVARD RD ASHEVILLE	164		0.097	0.095	0.094	0.090	0.087	0.086	7
37-027-0003 CALDWELL	HWY 321 NORTH LENOIR	206		0.092	0.090	0.089	0.085	0.084	0.083	4
37-029-0099 CAMDEN	COUNTY ROAD 1136 & 1134	198		0.084	0.076	0.073	0.072	0.071	0.071	0
37-033-0001 CASWELL	CHERRY GROVE RECREATION	213		0.102	0.102	0.093	0.092	0.092	0.089	9
37-037-0004 CHATHAM	RT 4 BOX 64 PITTSBORO NC 27312	213		0.084	0.083	0.082	0.082	0.081	0.079	0
37-051-0008 CUMBERLAND	1/4MI SR1857/US301/1857	211		0.093	0.092	0.089	0.087	0.084	0.084	4
37-051-1003 CUMBERLAND	3625 GOLFVIEW ROAD HOPE MILLS	203		0.098	0.093	0.093	0.084	0.084	0.082	3
37-059-0002 DAVIE	246 MAIN STREET COOLEEMEE	213		0.098	0.098	0.096	0.095	0.094	0.093	17
37-061-0002 DUPLIN	HWY 50 KENANSVILLE	212		0.092	0.085	0.084	0.081	0.081	0.079	2
37-063-0013 DURHAM	2700 NORTH DUKE STREET DURHAM	210		0.098	0.097	0.092	0.090	0.089	0.087	7
37-065-0099 EDGECOMBE	RT 2, BOX 195 TARBORO	201		0.096	0.095	0.086	0.085	0.084	0.084	4
37-067-0022 FORSYTH	1300 BLK. HATTIE AVENUE WINSTON-SALEM	178		0.094	0.094	0.092	0.090	0.086	0.086	6
37-067-0027 FORSYTH	7635 HOLLYBERRY LANE WINSTON-SALEM	177		0.091	0.082	0.082	0.080	0.079	0.079	1
37-067-0028 FORSYTH	6496 BAUX MOUNTAIN RD WINSTON-SALEM	175		0.094	0.091	0.088	0.086	0.085	0.082	5
37-067-1008 FORSYTH	3656 PIEDMONT MEMORIAL DRIVE WINSTON-SALEM	179		0.092	0.091	0.090	0.089	0.088	0.087	9
37-069-0001 FRANKLIN	431 S. HILLBOROUGH ST FRANKLINTON	206		0.097	0.091	0.089	0.088	0.085	0.082	5
37-077-0001 GRANVILLE	WATER TREATMENT PLANT JOHN UMSTEAD HOSPITAL BUTNER	213		0.093	0.091	0.091	0.091	0.090	0.088	10

SITE NUMBER	ADDRESS	NUM	VALID DAILY 8-HOUR MAXIMUM						NO. VALUES >.085 MEAS
			COUNTY	OBS	1ST	2nd	3rd	4th	
37-081-0011 GUILFORD	KEELY PARK, KEELY RD, MCCLEANSVILLE	211	0.099	0.096	0.094	0.090	0.089	0.088	8
37-087-0004 HAYWOOD	2177 SHEVILLE ROAD WAYNESVILLE	143	0.093	0.088	0.086	0.083	0.082	0.082	3
37-087-0035 HAYWOOD	TOWER BLUE RIDGE PARKWAY MILE MARKER 410	85	0.091	0.089	0.087	0.085	0.081	0.079	4
37-087-0036 HAYWOOD	GREAT SMOKY MOUNTAIN NATIONAL PARK	144	0.095	0.089	0.087	0.087	0.085	0.082	5
37-099-0005 JACKSON	BARNET KNOB FIRE TOWER	100	0.095	0.091	0.086	0.085	0.083	0.080	4
37-101-0002 JOHNSTON	3411 JACK ROAD CLAYTON	211	0.105	0.094	0.085	0.082	0.081	0.080	3
37-107-0004 LENOIR	CORNER HWY 70 EAST KINSTON	209	0.098	0.090	0.084	0.081	0.081	0.081	2
37-109-0004 LINCOLN	RIVERVIEW ROAD LINCOLNTON	212	0.104	0.097	0.095	0.094	0.094	0.092	15
37-117-0001 MARTIN	HAYES STREET (#2WELL SITE)	214	0.089	0.085	0.084	0.084	0.084	0.083	2
37-119-0041 MECKLENBURG	1120 EASTWAY DRIVE CHARLOTTE	203	0.126	0.107	0.099	0.096	0.093	0.092	12
37-119-1005 MECKLENBURG	400 WESTINGHOUSE BLVD. CHARLOTTE	204	0.097	0.091	0.088	0.087	0.086	0.084	12
37-119-1009 MECKLENBURG	29 N@ MECKLENBURG CAB CO CHARLOTTE	203	0.112	0.110	0.107	0.101	0.101	0.100	16
37-129-0002 NEW HANOVER	6028 HOLLY SHELTER RD	204	0.099	0.088	0.084	0.081	0.079	0.078	2
37-131-0002 NORTHAMPTON	ROUTE 46 GASTON	210	0.086	0.083	0.080	0.080	0.080	0.076	1
37-145-0099 PERSON	SR 1102 & NC 49	212	0.088	0.084	0.084	0.082	0.082	0.082	1
37-147-0099 PITT	US 264 NEAR WATER TOWER FARMVILLE	210	0.110	0.096	0.086	0.083	0.081	0.080	3
37-157-0099 ROCKINGHAM	6371 NC 65 @ BETHANY SCHOOL BETHANY	213	0.098	0.091	0.086	0.082	0.079	0.077	3
37-159-0021 ROWAN	WEST ST & GOLD HILL AVENUE ROCKWELL	214	0.103	0.102	0.094	0.092	0.089	0.086	7
37-159-0022 ROWAN	925 N ENOCHVILLE AVE ENOCHVILLE	203	0.105	0.096	0.095	0.095	0.094	0.094	12
37-173-0002 SWAIN	CENTER STREET PARKS 7 REC FACILITY	197	0.076	0.075	0.075	0.074	0.070	0.069	0
37-179-0003 UNION	701 CHARLES STREET MONROE	211	0.088	0.087	0.086	0.085	0.084	0.081	4

SITE NUMBER COUNTY	ADDRESS	NUM OBS	VALID DAILY 8-HOUR MAXIMUM						NO. VALUES >.085 MEAS
			1ST	2nd	3rd	4th	5TH	6TH	
37-183-0014 WAKE	E. MILLBROOK JR HI 3801 SPRING FOREST ROAD RALEIGH	212	0.097	0.091	0.088	0.087	0.086	0.086	6
37-183-0015 WAKE	808 NORTH STATE STREET RALEIGH	211	0.099	0.097	0.096	0.089	0.088	0.085	6
37-183-0016 WAKE	201 NORTH BROAD STREET FUQUAY-VARINA	211	0.100	0.087	0.087	0.085	0.085	0.085	6
37-183-0017 WAKE	5033 TV TOWER ROAD GARNER	207	0.107	0.102	0.089	0.082	0.081	0.081	3
37-199-0003 YANCY	BLUE RIDGE PARKWAY	181	0.093	0.088	0.088	0.087	0.084	0.082	4
Total Samples		8,841							239
Total Sites Sampled		45							




Figure 5.11 Ozone: Second Highest Annual 1-Hour Average, 2000




Figure 5.12 Ozone: Second Highest Annual 8-Hour Average, 2000




Figure 5.13 Distribution of Statewide Fourth-Maximum 8-Hour Ozone Concentrations, 1991-2000, and Smoothed Regression Trend Line




Figure 5.14 Number of Exceedances of the Daily Maximum 8-Hour Ozone Averages of 0.085 ppm or Greater, 1993-2000




Figure 5.15 Number of Exceedances of the Daily Maximum 8-Hour Ozone Averages of 0.085 ppm or Greater, by Region, 1993-2000.

Vertical bars represent (left to right)
 Charlotte-Gastonia, Greensboro-Winston-Salem-High Point, Raleigh-Durham, and all other areas.

5.6 Sulfur Dioxide

Sulfur dioxide (SO₂) concentrations were measured by the State and two local program agencies using EPA reference or equivalent methods. Eight SO₂ monitors were active in North Carolina in 2000. Some SO₂ sites are operated only every third year. Ten sites provided data in 1999, (and will next be operated in 2002), six sites provided data in 1998 (and will next be operated in 2001).

From the 24 sites with SO₂ data obtained between 1998 and 2000, 143,048 valid hourly averages were collected. A map of the active SO₂ sampling sites is presented in Figure 5.16 and a detailed summary of the data from each site is given in Table 5.7.

There were no exceedances of the SO₂ ambient air quality standards in 2000. The highest annual arithmetic mean was 27 $\mu\text{g}/\text{m}^3$, or about 33 percent of the standard

(80 $\mu\text{g}/\text{m}^3$). The highest maximum 24-hour average was 81 $\mu\text{g}/\text{m}^3$, about 22 percent of the standard (365 $\mu\text{g}/\text{m}^3$), and the highest maximum 3-hour average was 325 $\mu\text{g}/\text{m}^3$, about 25 percent of the welfare-related (secondary) standard.

Apparently, the size of an urban area has little effect on the ambient concentrations of SO₂ in North Carolina. Seasonal variations, such as those with CO and O₃, do not appear to exist for SO₂. Major source characteristics such as type, size, distribution, control devices, operating conditions and dispersion situations significantly affect the amount of SO₂ in ambient air.

The second highest three-hour concentrations in each county are shown in Figure 5.17. The second highest 24-hour concentrations in each county are shown in Figure 5.18.




Figure 5.16 Location of Sulfur Dioxide Monitoring Sites

Table 5.7 Sulfur Dioxide in Parts Per Million from All Sites for 1998-2000

SITE NUMBER COUNTY	ADDRESS	NUM OBS	ONE-HOUR MAXIMA		THREE-HOUR MAXIMA		24-HOUR MAXIMA		ARITH MEAN
			1ST	2ND	1ST	2ND	1ST	2ND	
2000 Data									
37-013-0003 BEAUFORT	NC HIGHWAY 306	6,671	0.178	0.085	0.107	0.057	0.032	0.020	0.004
37-059-0002 DAVIE	246 MAIN STREET COOLEEMEE	8,249	0.071	0.054	0.055	0.049	0.019	0.018	0.004
37-067-0022 FORSYTH	1300 BLK. HATTIE AVE WINSTON-SALEM	8,420	0.086	0.078	0.057	0.057	0.020	0.019	0.005
37-109-0004 LINCOLN	RIVERVIEW ROAD LINCOLNTON	8,159	0.076	0.066	0.046	0.043	0.022	0.018	0.004
37-119-0041 MECKLENBURG	1120 EASTWAY DRIVE CHARLOTTE	8,539	0.100	0.075	0.059	0.045	0.017	0.017	0.004
37-129-0006 NEW HANOVER	HWY 421 NORTH WILMINGTON	8,304	0.088	0.084	0.062	0.061	0.039	0.030	0.006
37-131-0002 NORTHAMPTON	RT 46 GASTON	8,361	0.037	0.033	0.025	0.022	0.012	0.012	0.004
37-147-0099 PITT	US 264 NEAR WATER TWR FARMVILLE	8,153	0.023	0.023	0.018	0.016	0.009	0.007	0.003
Total Samples		64,856							
Total Sites Sampled		8							
1999 Data									
37-013-0003 BEAUFORT	NC HIGHWAY 306	7,145	0.071	0.062	0.047	0.040	0.019	0.015	0.006
37-013-0004 BEAUFORT	SOUTH FERRY LANDING PAMLICO RIVER	5,181	0.063	0.060	0.043	0.037	0.017	0.015	0.005
37-051-1003 CUMBERLAND	3625 GOLFVIEW ROAD HOPE MILLS	8,271	0.018	0.018	0.012	0.012	0.007	0.007	0.005
37-061-0002 DUPLIN	HWY 50 KENANSVILLE	8,258	0.017	0.016	0.014	0.013	0.007	0.007	0.005
37-065-0099 EDGECOMBE	RT 2, BOX 195 TARBORO	7,416	0.063	0.060	0.012	0.011	0.008	0.008	0.005
37-101-0002 JOHNSTON	3411 JACK ROAD CLAYTON	7,567	0.042	0.037	0.033	0.019	0.013	0.009	0.005
37-119-0034 MECKLENBURG	PLAZA RD. & LAKEDELL CHARLOTTE	7,221	0.083	0.082	0.066	0.047	0.014	0.013	0.004
37-129-0006 NEW HANOVER	HWY 421 NORTH WILMINGTON	8,280	0.347	0.160	0.125	0.072	0.031	0.027	0.009
Total Samples		73,118							
Total Sites Sampled		10							

1998 Data										
37-037-0004 CHATHAM	RT4 BOX62 PITTSBORO	8,283	0.050	0.036	0.030	0.028	0.013	0.009	0.005	
37-117-0001 MARTIN	HAYES STREET (#2WELL SITE)	8,184	0.014	0.013	0.012	0.012	0.007	0.006	0.005	
37-145-0099 PERSON	SR 1102 & NC49	8,265	0.077	0.073	0.067	0.056	0.021	0.016	0.006	
37-159-0021 ROWAN	WEST STREET & GOLD HILL AVENUE	2,891	0.065	0.061	0.036	0.031	0.012	0.010	0.006	
37-159-OO22 ROWAN	ROCKWELL 925 N ENOCHVILLE AVE	2,712	0.087	0.074	0.067	0.036	0.016	0.012	0.006	
37-173-0002 SWAIN	CENTER ST PARKS 7 REC FACILITY	7,869	0.075	0.019	0.028	0.018	0.008	0.008	0.005	
Total Samples		38,204								
Total Sites Sampled		6								




Figure 5.17 Sulfur Dioxide: Second Highest 3-Hour Averages in the Most Recent Year of Data from 1998, 1999 or 2000




Figure 5.18 Sulfur Dioxide: Second Highest 24-Hour Averages in the Most Recent Year of Data from 1998, 1999 or 2000

5.7 Nitrogen Dioxide

Nitrogen dioxide (NO₂) concentrations were measured using EPA reference or equivalent continuous monitors in 2000 at one local program site in Forsyth County and one local program site in Mecklenburg County.

From these two sites, 12,688 hourly NO₂ measurements were reported. A map of the

NO₂ sampling sites is presented in Figure 5.19, and a summary of the 2000 NO₂ data is given in Table 5.8.

Each urban area site has only a few outlying high hourly sample values that are above the standard defined for the annual arithmetic mean. The arithmetic means (Table 5.8) are about 34 percent of the standard.




Figure 5.19 Location of Nitrogen Dioxide Monitoring Sites

Table 5.8 Nitrogen Dioxide in Parts Per Million for 2000

SITE NUMBER COUNTY	ADDRESS	NUM OBS	ONE-HOUR MAXIMA		ARITH MEAN
			1ST	2ND	
37-067-0022 FORSYTH	1300 BLK. HATTIE AVENUE WINSTON-SALEM	4,033	0.082	0.079	0.018
37-067-0041 MECKLENBURG	1120 EASTWAY DRIVE CHARLOTTE	8,655	0.094	0.088	0.018
Total Samples		12,688			
Total Sites Sampled		2			

5.8 Lead

The state and local program agencies have not performed routine analysis of ambient lead (Pb) in North Carolina since 1982. Lead monitoring was discontinued as a result of the low measurements and a continuing decrease in the lead concentrations being reported. The decrease in ambient Pb concentrations is due to the reduction and elimination of leaded gasoline, resulting in greatly reduced lead emissions from automobiles.

The most recent year of data available prior to 1996-97 was in 1990. Because the previous data was so old, the state began metals analysis at three locations on 1996. These metal sites will be relocated to other locations in future years. The purpose of these sites is to gather background information about lead and other metals. No lead sites operated in 2000.

The change in analytical laboratories from the EPA's National Particulate Analysis Program to the state program also changed the minimum detectable levels of the method from 0.01 to 0.04 $\mu\text{g}/\text{m}^3$, respectively. Concentrations of most metals are below detectable limits regardless of the method used.

During 1999, a special study focusing on arsenic levels was undertaken. Lead, as well as 9 other toxic metals were analyzed on TSP samplers with PM_{10} filters at 10 ambient air monitoring sites, by a contract laboratory using inductively coupled plasma/mass spectrometry (ICP/MS). This method has a sensitivity of three orders of magnitude greater than the past methods. Of the 526 samples analyzed only 39 exceeded the minimum detectable level of 0.01 $\mu\text{g}/\text{m}^3$ and only 2 exceeded the minimum detectable level of 0.04 $\mu\text{g}/\text{m}^3$, with values of 0.04154 $\mu\text{g}/\text{m}^3$ and 0.04174 $\mu\text{g}/\text{m}^3$. This study was discontinued in 2000. The lead standard remains 1.5 $\mu\text{g}/\text{m}^3$ for a quarterly average.

6 Air Quality Index

The Air Quality Index (AQI) was developed by the EPA to provide the public with a simple, accessible, and uniform assessment of air quality at a specific location, based on the criteria pollutants PM_{2.5}, PM₁₀, CO, O₃ (both 1 and 8 hour values), SO₂ and NO₂. AQI measurements are made and reported in all U.S. metropolitan statistical areas (MSA) with a population over 350,000. Ambient concentrations for each of these seven pollutants are converted to a numerical scale ranging from 0 to 500, where 100 corresponds to the EPA primary standard for a 24-hour average (8-hour CO average, 1 and 8-hour O₃ average) and 500 corresponds to a concentration associated with significant harm. The AQI is determined by the pollutant with the highest scaled concentration, and a subjective description of "Good", "Moderate", "Unhealthy for sensitive groups", "Unhealthy", "Very unhealthy", or "Hazardous" is included with the report, with the descriptions corresponding to AQI values of 0-50, 51-100, 101-150, 151-200, 201-300, and 301-500, respectively. For AQI values between 101 and 500, an appropriate cautionary statement is included advising people susceptible to deleterious health effects to

restrict activities and exposure to the ambient air.

An AQI of 101-200 (unhealthy for sensitive groups and unhealthy) can produce mild aggravation of symptoms in susceptible persons and possible irritation in healthy persons. People with existing heart or lung ailments should reduce physical exertion and outdoor activity. The general population should reduce vigorous outdoor activity.

An AQI of 201 to 300 (very unhealthy) can produce significant aggravation of symptoms and decreased exercise tolerance in persons with heart or lung disease, and a variety of symptoms in healthy persons. Elderly people and those with existing heart or lung disease should stay indoors and reduce physical activity. The general population should avoid vigorous outdoor activity.

The health effects of an AQI of over 300 (hazardous) include early onset of certain diseases in addition to significant aggravation of symptoms and decreased exercise tolerance in healthy persons. The elderly and persons with existing diseases

should stay indoors and avoid physical exertion.

At AQI values over 400, premature death of ill and elderly persons may result, and healthy people will experience adverse symptoms that affect normal activity. Outdoor activity should be avoided. All people should remain indoors, keeping windows and doors closed, and should minimize physical exertion.

During winter months in North Carolina, carbon monoxide usually has the highest air quality index value, and in summer months

the highest index value is usually due to ozone.

In 2000, nine areas provided an AQI report to the public by telephone using computer-generated recorded voice announcements 24 hours daily. These areas are identified in the box below. The AQI report also may be published by local newspapers or broadcast on radio and television stations.

The Air Quality Index report is now available by telephone for nine areas, as follows:

Statewide toll-free
(for Asheville, Durham, Fayetteville, Greensboro, Greenville,
Raleigh, Wilmington, and Winston-Salem areas)

888-AIR-WISE

Charlotte area

704-333-SMOG

In 2002, we will provide an AQI Report on the North Carolina DAQ web site, (<http://www.daq.state.nc.us/monitor>).

Air Quality Index values during 2000 at six metropolitan areas in North Carolina are given in Figures 6.1, 6.2, 6.3, 6.4, 6.5 and 6.6.




Figure 6.1 Daily Air Quality Index Values for Asheville, NC, Metropolitan Statistical Area, 2000




Figure 6.2 Daily Air Quality Values for Charlotte-Gastonia, NC,-Rock Hill, SC, Metropolitan Statistical Area, 2000




Figure 6.3 Daily Air Quality Index Values for Fayetteville, NC, Metropolitan Statistical Area, 2000




Figure 6.4 Daily Air Quality Index Values Greensboro-Winston-Salem-High Point, NC, Metropolitan Statistical Area, 2000




Figure 6.5 Daily Air Quality Index Values for Raleigh-Durham, NC, Metropolitan Statistical Area, 2000




Figure 6.6 Daily Air Quality Index Values for Wilmington, NC, Metropolitan Statistical Area, 2000

7 Acid Rain

7.1 Sources

Acid rain is produced when nitrate and sulfate ions from automobile and industrial sources are released into the atmosphere, undergo a reaction with moisture in the air, and are deposited as acid precipitation. Acid ions are produced when sulfur dioxide and nitrogen oxides reach equilibrium with water to form sulfuric acid and nitric acid.

7.2 Effects

Many agricultural crops in North Carolina are sensitive to acid rain. Forests are subject to mineral loss from acid rain exposure and may also suffer root damage. Acid fogs and mists, typical in the mountains of North Carolina, can expose trees and plants to even higher acid concentrations and cause direct damage to foliage. Lakes, rivers and streams that are too acidic can impede fish and plant growth.

7.3 Monitoring

Acid rain monitoring has been conducted nationally, including in North Carolina, since 1978 by the National Atmospheric Deposition Program (NADP) and the National Trends Network (NTN) which merged with NADP in 1982. In 2000, acid rain samples were collected at seven sites in North Carolina and one Tennessee site in the Great Smoky Mountains less than 10 miles from the western border of North Carolina.

NADP conducts acid deposition monitoring using a wet/dry bucket type sampler. When rainfall is detected, a sensor is activated and a metal lid automatically covers and protects the Adry≡ sample, exposing the Awet≡ bucket to collect precipitation.

Acidity is measured using a ApH≡ scale. The pH scale is numbered from 0 to 14, with 0 being extremely acidic and 14 being extremely basic. A substance with a pH of five is ten times as acidic as one with a pH of six, 100 times as acidic as a substance with a pH of seven, etc. Neutral water with an equal concentration of acid and base ions has a pH of seven. The pH of vinegar is approximately 2.8, and lemon juice has a pH of about 2.3. The pH of ammonia is approximately 12.

Pure water in equilibrium with the air is slightly acidic and has a pH of approximately 5.6. The measurements of pH at the North Carolina monitoring sites in 2000 ranged from 4.39 to 4.73 with a mean of 4.59. The 2000 pH annual means for North Carolina from the NADP database are presented in Figure 7.1 and Table 7.1. Table 7.1 also exhibits conductivity averages and precipitation totals for rainfall. Measured concentrations of several other chemical constituents of precipitation are given in Table 7.2.

The highest pH (and the least acid) precipitation occurred at the Sampson County site. This general area in southeastern North Carolina has the greatest numbers of animal producing farms. This area has the highest emissions of ammonia, a basic gas emitted from animal wastes. Table 7.2 shows that the ammonium concentration in precipitation is highest at the Sampson County site.




Figure 7.1 Annual Mean pH Values at North Carolina NADP Sites, 2000

Table 7.1 pH, Conductivity in Microsiemens per Centimeter and Precipitation in Inches from the National Atmospheric Deposition Program for 2000.

COUNTY SITE ADDRESS	pH	CONDUCTIVITY	PRECIPITATION
BERTIE 340320 LEWISTON	4.62	15.4	39.3
MACON 342500 COWEETA	4.64	13.4	42.9
ROWAN 343460 PIEDMONT RESEARCH STATION	4.39	23.8	30.1
SAMPSON 343560 CLINTON CROPS RESEARCH STATION	4.73	14.9	43.3
SCOTLAND 343600 JORDAN CREEK	4.48	19.1	41.9
WAKE 344160 FINLEY FARM	4.56	17.3	42.5
YANCEY 344500 Mt. MITCHELL	4.67	12.2	66.2
SEVIER (TN) 441190 GREAT SMOKY MTS NATIONAL PARK ELKMONT TN	4.64	14.1	56.0

Table 7.2 Ion Concentrations in Milligrams per Liter (Precipitation-weighted Annual Means) from the National Atmospheric Deposition Program Data for 2000.

COUNTY SITE	% COMPLETE-NESS	Ca	Mg	K	Na	NH4	NO3	CL	SO4
BERTIE 340320	89.9	0.07	0.025	0.024	0.175	0.23	1.01	0.31	1.28
MACON 342500	85.8	0.07	0.012	0.016	0.074	0.15	0.81	0.13	1.14
ROWAN 343460	90.2	0.09	0.019	0.056	0.082	0.32	1.37	0.21	2.14
SAMPSON 343560	89.6	0.08	0.026	0.028	0.192	0.46	1.13	0.34	1.50
SCOTLAND 343600	87.7	0.07	0.020	0.021	0.135	0.21	1.16	0.25	1.55
WAKE 344160	91.6	0.06	0.017	0.018	0.122	0.32	1.12	0.23	1.55
YANCEY 344500	77.7	0.06	0.008	0.014	0.039	0.16	0.66	0.07	1.11
SEVIER (TN) 441190	85.8	0.08	0.014	0.050	0.055	0.19	0.90	0.11	1.21

References

- Cornelius, Wayne L. (1996). Effects of North Carolina's Oxygenated Fuel Program on Ambient Carbon Monoxide Concentrations. *Air Quality Section, Division of Environmental Management, N.C. Dept. Of Env., Health, and Nat. Res.*
- Cornelius, Wayne L. (1997). Comparison of Nitrogenous Ion Deposition and Human and Animal Census Trends in Eastern North Carolina. *Air Quality Section, Division of Environmental Management, N.C. Dept. Of Env., Health, and Nat. Res.*
- Maynard, Andrew D. and Paul a. Jensen (2001). Aerosol Measurement in the Workplace. Chapter 25 in *Aerosol Measurement: Principles, Techniques, and Applications, Second Edition*, Paul A. Baron and Klaus Willeke, eds., p. 779-799.
- NADP (National Atmospheric Deposition Program) (1995). Notification of Important Change in NADP/NTN Procedures on 11 January 1994.
- North Carolina Department of Environment, Health, and Natural Resources (1991a). 1989 Ambient Air Quality Report. *Air Quality Section, Division of Environmental Management, N.C. Dept. Of Env., Health, and Nat. Res.*
- North Carolina Department of Environment, Health, and Natural Resources (1991b). Ambient Air Quality Trends in North Carolina 1972-1989. *Air Quality Section, Division of Environmental Management, N.C. Dept. Of Env., Health, and Nat. Res.*
- United States Environmental Protection Agency [US EPA] (1993), A Code of Federal Regulations, Title 40, Parts 1 to 51, Protection of Environment, (July 1 ed.). *Office of the Federal Register (National Archives and Records Administration)*, Washington, DC.
- Watson, John G. and Judith C. Chow (2001). Ambient Air Sampling. Chapter 27 in *Aerosol Measurement: Principles, Techniques, and Applications, Second Edition*, Paul A. Baron and Klaus Willeke, eds., p. 821-844.

Appendix A. Air Pollution Monitoring Agencies

North Carolina State Headquarters

Division of Air Quality

Parker Lincoln Building
2728 Capital Boulevard
1641 Mail Service Center
Raleigh, North Carolina 27699-1641
(919) 715-0665

North Carolina Regional Offices

Asheville Regional Office

Interchange Building
59 Woodfin Place
Asheville, North Carolina 28801
(828) 251-6208

Counties of Avery, Burke, Caldwell, Cherokee, Clay, Graham, Haywood, Henderson, Jackson, Macon, Madison, McDowell, Mitchell, Polk, Rutherford, Swain, Transylvania, and Yancey.

Fayetteville Regional Office

Suite 714
225 Green Street
Fayetteville, North Carolina 28301
(910) 486-1541

Counties of Anson, Bladen, Cumberland, Harnett, Hoke, Montgomery, Moore, Robeson, Richmond, Sampson, and Scotland.

Mooresville Regional Office

919 North Main Street
Mooresville, North Carolina 28115
(704) 663-1699

Counties of Alexander, Cabarrus, Catawba, Cleveland, Gaston, Iredell, Lincoln, Rowan, Stanly and Union.

Raleigh Regional Office

3800 Barrett Drive
Raleigh, North Carolina 27611
(919) 571-4700

Counties of Chatham, Durham, Edgecombe, Franklin, Granville, Halifax, Johnston, Lee, Nash, Northampton, Orange, Person, Vance, Wake, Warren, and Wilson.

Washington Regional Office

943 Washington Square Mall
Washington, North Carolina 27889
(252) 946-6481

Counties of Beaufort, Bertie, Camden, Chowan, Craven, Currituck, Dare, Gates, Greene, Hertford, Hyde, Jones, Lenoir, Martin, Pamlico, Pasquotank, Perquimans, Pitt, Tyrrell, Washington, and Wayne.

Wilmington Regional Office

127 Cardinal Drive Extension
Wilmington, North Carolina 28405-3845
(910) 395-3900

Counties of Brunswick, Carteret, Columbus, Duplin, New Hanover, Onslow and Pender.

Winston-Salem Regional Office

585 Waughtown Street
Winston-Salem, North Carolina 27107
(336) 771-4600

Counties of Alamance, Alleghany, Ashe, Caswell, Davidson, Davie, Guilford, Rockingham, Randolph, Stokes, Surry, Yadkin, Watauga, and Wilkes.

Local Agencies in North Carolina

Forsyth County Environmental Affairs Department

537 North Spruce Street
Winston-Salem, North Carolina 27101
(336) 727-8064

Mecklenburg County Department of Environmental Protection

700 N. Tryon Street, Suite 205
Charlotte, North Carolina 28202-2236
(704) 336-5500

Western North Carolina Regional Air Pollution Quality Agency (Buncombe County and Asheville)

49 Mount Carmel Road
Asheville, North Carolina 28806
(828) 255-5655

Tribal Agency in North Carolina

Eastern Band of Cherokee Indians

Tribal Environmental Office
P. O. Box 455
Cherokee, North Carolina 28719
(828) 497-3814


Appendix B. Exceptional Events

Type of Event	Pollutants Affected
Natural Events	
Sustained high wind speeds	particulate matter (PM)
Stagnations, inversions	all pollutants
Unusual lack of precipitation	PM
Stratospheric ozone intrusion	O ₃
Volcanic eruption	CO, SO ₂ , PM
Forest fires	CO, PM, O ₃
High pollen count	PM
Unintentional Man-made Events	
Large structural fires	CO, PM
Major traffic congestion due to accident or nonrecurring obstruction	CO
Chemical spills	SO ₂ , NO ₂ , PM, CO
Industrial accidents	SO ₂ , NO ₂ , PM, CO
Intentional Man-made Events	
Short-term construction/demolition	PM
Sandblasting	PM
High-sulfur oil refining	SO ₂
Roofing operations	PM, SO ₂
Salting or sanding of streets	PM
Infrequent large gatherings	PM, CO
Soot blowing from ships	PM
Agricultural tilling	PM
Prescribed burning	CO, PM
Noncompliance of local sources	CO, SO ₂

Appendix C. Box-And-Whisker Plots

A *box-and-whisker plot* (also called *boxplot* or *schematic plot*) is a schematic diagram useful for depicting the location, spread and skewness of a continuous data variable. Box plots are constructed from *order statistics* (data values sorted from smallest to largest). The "box" of the box plot is oriented parallel to a continuous scale and is defined by 3 points, (1) a line or point in the interior of the box at the median of the data (a point that divides the order statistics into two equal parts), and (2) upper and (3) lower *fourths* or *quartiles*. (Fourths divide the upper and lower halves of the data values into two equal parts; quartiles divide the entire range of the data into 4 equal parts. Fourths and quartiles are not necessarily the *same*, because there may be more than one number that appropriately divides a given set of data in the prescribed way, and different computational techniques [or computer programs] may make different choices.)

The distance between the upper and lower fourth in the box plot is called the *interquartile range*. In most box plots, the length of each of the *whiskers* is 1.5 times the interquartile range or to the extreme (maximum or minimum) of the data, whichever is *shorter*. The endpoint of each whisker is called an *inner fence*. (In the box plots pictured below, the end of each whisker is marked by a "staple" for clarity.) There may be data points, called *outliers*, beyond the inner fences; if so, they are usually indicated individually on the box plot by a dot, small circle, or (as below) a short line segment perpendicular to the axis of the box. Box plots of variables with very



long-tailed distributions may display two kinds of outliers: small dots for those just beyond the inner fences and larger dots or circles for *extreme outliers* at a distance of more than 3.0 times the interquartile range beyond the fourths. This boundary between outliers and extreme outliers is termed the *outer fence* and usually not explicitly shown in the plot.

The maximum and minimum values are always visible in a box-and-whisker plot as either the outermost outliers or, if there is no outlier, the position of the inner fence.

Box plots may have additional, optional features, such as a point marker at the *arithmetic mean* or a distinctive display of a *confidence interval for the median*, which is calculated from the fourths. In the figure, the arithmetic mean is marked with an "X", and the confidence interval for the median is displayed as a shaded or colored range; it is also common to display the confidence interval by cutting notches in the sides of the box at its endpoints.

Box plots are very useful for comparing two or more variables by placing two comparable variables side-by-side on the same scale (as in the figure). The statistics displayed can be directly compared, and statistical significance of difference between the medians can be assessed by examining overlap or lack of overlap of confidence intervals.

Appendix D. Nonattainment and North Carolina

What is nonattainment and what are the sources of the pollutants?

The United States Environmental Protection Agency (EPA) sets National Ambient Air Quality Standards. North Carolina monitors concentrations of air pollutants in the ambient air. Some of these monitors have measured concentrations of ozone and carbon monoxide exceeding the standards. Areas that have not met the National Ambient Air Quality Standards can be classified by EPA as Nonattainment.≡

Mobile sources such as cars and trucks are the primary cause of carbon monoxide and ozone precursors. About 90 percent of the carbon monoxide emissions come from motor vehicles. In the urban areas, 60 percent of the nitrogen oxides and 25 percent of the man-made hydrocarbons or volatile organic compound emissions come from motor vehicles; the rest comes from off-road vehicles, utility and industrial boilers, petroleum marketing, factories, businesses, and households. Nitrogen oxides react with volatile organic compounds and sunlight in warm weather to produce ozone.

Why is my county nonattainment?

EPA guidance recommends that an entire Metropolitan Statistical Area (MSA) be designated nonattainment when a monitor is found to be violating the National Ambient Air Quality Standards (NAAQS). This policy is due to the regional nature of certain pollutants, like ozone. Ozone is formed in the atmosphere under complex chemical reactions. Sometimes the ozone levels are higher just downwind of urban areas because of the time it takes the pollutants to react to form ozone. Therefore, larger areas are designated nonattainment to represent the likely area contributing to the air quality problems.

Once we are nonattainment, what is the process for becoming attainment?

North Carolina is required by the federal Clean Air Act and EPA to produce and implement emission reduction plans and show that these plans are strong enough to produce compliance with the standards. The plans could involve resource-intensive monitoring, emissions inventory, modeling, public participation, and strategy formulation efforts. There are deadlines for producing the plans and for achieving compliance with the standards. EPA must approve the plans.

How does the public get involved in the formulation of the emission reduction plans, known as State Implementation Plan (SIP) revisions?

Local agencies and officials, as well as state agencies, will be involved in drawing up the SIP revisions. There will be public meetings or special citizen panels. When draft SIP revisions are done, there will be public hearings on them. The SIP revisions must be approved by the N.C.

Environmental Management Commission and possibly by local bodies as well. The N. C. General Assembly also reviews the SIP. EPA's approval process also includes an opportunity for public comment.

How will it affect citizens?

Emission reduction strategies fall into several categories. Motor vehicle inspection/maintenance may be required for hydrocarbons, carbon monoxide, and nitrogen oxides. Traffic patterns may be altered by changing roads or traffic signals. Both new and existing factories and business may have to reduce emissions by installing control equipment or changing processes. Cleaner burning gasoline may be required. More controls will be required on utility and industrial boilers. All of these measures may mean higher costs to the public.

What happens if North Carolina refuses to address these air pollution problems?

Under the Clean Air Act, EPA has the authority to apply sanctions. EPA can require more stringent offsets for new facilities of major pollutant sources, and may withhold federal highway construction funds in the nonattainment areas.

What is the likelihood of receiving sanctions if we are showing progress in reducing pollution?

North Carolina can avoid sanctions if it produces and carries out SIP revisions that EPA approves by the deadlines. If pollution concentrations do not recede and attain the standards as projected, the EPA could impose construction bans. However, EPA has some discretion about imposing sanctions. Sanctions are a last step to persuade states to take required positive action.

What does inspection/maintenance cost?

As of December, 2001, the inspection/maintenance (I/M), or motor vehicle tailpipe testing process, costs the motorist \$30.00. If a vehicle fails the test, it must be repaired. A waiver is available if a vehicle still fails after \$250.00 worth of repairs have been done. The \$250.00 limit does not apply to tampered or misfueled vehicles. The inspection/maintenance program includes tests for hydrocarbon (HC) and carbon monoxide (CO) emissions for the following counties, Mecklenburg, Wake, Guilford, Forsyth, Durham, Gaston, Cabarrus, Orange, and Union. The number of counties will increase to 48 by 2006 under the Clean Air bill passed in 2000. The cost for this new test will be set by the General Assembly during the 2001 legislative session. Currently, only gasoline powered motor vehicles built after 1974, excluding the current model year and motorcycles, are inspected in these counties. Inspection/maintenance pass/fail levels vary with vehicle age and pollutant.