

2010 USE SUPPORT & METHODOLOGY

IN THE ROANOKE RIVER BASIN

2010 IR CATEGORY	INTEGRATED REPORTING CATEGORIES FOR INDIVIDUAL ASSESSMENT UNIT/USE SUPPORT CATEGORY/PARAMETER ASSESSMENTS. A SINGLE AU CAN HAVE MULTIPLE ASSESSMENTS DEPENDING ON DATA AVAILABLE AND CLASSIFIED USES.
1	All designated uses are monitored and supporting
1b	Designated use was impaired, other management strategy in place and no standards violations for the parameter of interest (POI)
1nc	DWQ have made field determination that parameter in exceedance is due to natural conditions
1r	Assessed as supporting watershed is in restoration effort status
1t	No criteria exceeded but approved TMDL for parameter of interest
2	Some designated uses are monitored and supporting none are impaired Overall only
2b	Designated use was impaired other management strategy in place and no standards violations Overall only
2r	Assessed as supporting watershed is in restoration effort status overall only
2t	No criteria exceeded but approved TMDL for POI Overall only
3a	Instream/monitoring data are inconclusive (DI)
3b	No Data available for assessment
3c	No data or information to make assessment
3n1	Chlorophyll a exceeds TL value and SAC is met-draft
3n2	Chlorophyll a exceeds EL value and SAC is not met first priority for further monitoring-draft
3n3	Chlorophyll a exceeds threshold value and SAC is not met first second priority for further monitoring-draft
3n4	Chlorophyll a not available determine need to collect-draft
3t	No Data available for assessment –AU is in a watershed with an approved TMDL
4b	Designated use impaired other management strategy expected to address impairment
4c	Designated use impaired by something other than pollutant
4cr	Recreation use impaired no instream monitoring data or screening criteria exceeded
4cs	Shellfish harvesting impaired no instream monitoring data-no longer used
4ct	Designated use impaired but water is subject to approved TMDL or under TMDL development
4s	Impaired Aquatic Life with approved TMDL for Aquatic Life POI or category 5 listing
4t	Designated use impaired approved TMDL
5	Designated use impaired because of biological or ambient water quality standards violations and needing a TMDL
5r	Assessed as impaired watershed is in restoration effort status

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Grassy Creek-John H Kerr Reservoir Watershed		0301010208	
Roanoke River Basin		John H Kerr Reservoir-Roanoke River Subbasin		03010102	
Roanoke River Basin		Grassy Creek-John H Kerr Reservoir Watershed		0301010208	
⊙ 23-2-(1)	Grassy Creek (Grass Creek)	From source to John H. Kerr Reservoir at Granville County SR 1431		18.3 FW Miles	C
3a	Ecological/biological Integrity Benthos	Not Rated Bioclassification	Aquatic Life	2004	
⊙ 23-2-7-(1)	Johnson Creek	From source to Little Johnson Creek		5.3 FW Miles	C
1	Ecological/biological Integrity FishCom	Good-Fair Bioclassification	Aquatic Life	2004	
⊙ 23-2-3	Mountain Creek	From source to Grassy Creek		8.1 FW Miles	C
3a	Ecological/biological Integrity Benthos	Not Rated Bioclassification	Aquatic Life	2004	
⊙ 23-2-5	Rattlesnake Creek	From source to Grassy Creek		2.3 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
Roanoke River Basin		Butcher Creek-John H Kerr Reservoir Watershed		0301010209	
⊙ 23-4	Island Creek (Island Creek Reservoir)	From source to North Carolina-Virginia State Line, including that portion of Island Creek Reservoir in North Carolina below normal operating elevation		6.4 FW Miles	C
1	Ecological/biological Integrity Benthos	Good-Fair Bioclassification	Aquatic Life	2004	
⊙ 23-4-3	Little Island Creek (Vance County)	From source to Island Creek Reservoir, Island Creek		11.8 FW Miles	C
3a	Ecological/biological Integrity FishCom	Not Rated Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Nutbush Creek-John H Kerr Reservoir Watershed		0301010210	
⊙ 23-8-(1)a	Nutbush Creek (Including Nutbush Creek Arm of John H. Kerr Reservoir below normal pool elevation)	From source to NC 39		1.7 FW Miles	C
5	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	1994	1998

NC DWQ ROANOKE RIVER BASIN PLAN: 2010 USE SUPPORT 2011

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Nutbush Creek-John H Kerr Reservoir Watershed		0301010210	

⊙ **23-8-(1)b** **Nutbush Creek** From NC 39 to SR 1317 **1.6 FW Miles** **C**
(Including Nutbush Creek Arm of John H. Kerr Reservoir below normal pool elevation)

5	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	2004	1998
5	Ecological/biological Integrity FishCom	Fair Bioclassification	Aquatic Life	2004	1998
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	

⊙ **23-8-(2)** **Nutbush Creek Arm of John H. Kerr Reservoir (below normal pool elevation 300 feet MSL or as this elevation may be adjusted by the Corps of Engineers)** From Crooked Run to North Carolina-Virginia State Line **9,690.1 FW Acres** **B**

1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
---	--------------------------------------	----------------------	--------------	------	--

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Little Dan River-Dan River Watershed		0301010301	
Roanoke River Basin		Dan River Headwaters Subbasin		03010103	
Roanoke River Basin		Little Dan River-Dan River Watershed		0301010301	
⊙ 22-2	Archies Creek	North Carolina portion	7.3 FW Miles	C;Tr	
1	Ecological/biological Integrity FishCom	Excellent Bioclassification	Aquatic Life	2004	
⊙ 22-9	Big Creek	From source to Dan River	19.9 FW Miles	C;Tr	
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-12-(2)b	Cascade Creek	From dam at swimming lake to Dan River	4.3 FW Miles	B	
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
⊙ 22-12-(2)a	Cascade Creek (Hanging Rock Lake)	From backwaters to dam at swimming lake	12.2 FW Acres	B	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
⊙ 22-(1)a	DAN RIVER (North Carolina portion)	From North Carolina-Virginia State Line to Little Dan River	5.1 FW Miles	C;Tr	
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-(1)b	DAN RIVER (North Carolina portion)	From Little Dan River to Peters Creek	11.6 FW Miles	C;Tr	
1	Ecological/biological Integrity Benthos	Excellent Bioclassification	Aquatic Life	2004	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
3a	High Water Temperature	Potential Standards Violation	Aquatic Life	2008	
5	Turbidity	Standard Violation	Aquatic Life	2008	2008
⊙ 22-5	Elk Creek	From North Carolina-Virginia State Line to Dan River	2.9 FW Miles	C;Tr	
1	Ecological/biological Integrity FishCom	Good-Fair Bioclassification	Aquatic Life	2004	
⊙ 22-13-(2)	Indian Creek	From Window Falls to Dan River	2.7 FW Miles	C	
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
⊙ 22-18	Mill Creek	From source to Dan River	4.7 FW Miles	C	
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
⊙ 22-10	North Double Creek	From source to Dan River	14.0 FW Miles	C	
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
1	Ecological/biological Integrity FishCom	Good-Fair Bioclassification	Aquatic Life	2004	
⊙ 22-6	Peters Creek	From North Carolina-Virginia State Line to Dan River	9.1 FW Miles	C;Tr	
1	Ecological/biological Integrity FishCom	Excellent Bioclassification	Aquatic Life	2004	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Little Dan River-Dan River Watershed		0301010301	
⊙ 22-11	South Double Creek	From source to Dan River		9.9 FW Miles	B
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Town Fork Creek Watershed		0301010302	
⊙ 22-25-1	Brushy Fork Creek	From source to Town Fork Creek		3.0 FW Miles	C
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
⊙ 22-25a	Town Fork Creek	From source to Timmons Cr.		8.0 FW Miles	C
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
⊙ 22-25b	Town Fork Creek	From Timmons Cr. to Dan River		18.0 FW Miles	C
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Belews Lake-Dan River Watershed		0301010303	
⊙ 22-27-(7)	Belews Creek (including Belews Lake below elevation 725) (1)	From Southern Railroad Bridge to to a point 1.8 mile downstream of Forsyth-Stokes County Line		789.7 FW Acres	C
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
⊙ 22-27-(7.5)	Belews Creek (including Belews Lake below elevation 725) (1)	From a point 1.8 mile downstream of the Forsyth-Stokes County Line to Dan River, excluding the Arm of Belews Lake described below which are classified "WS-IV&B"		1,283.8 FW Acres	WS-IV
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-27-(1.5)	Belews Creek (Kernersville Lake)	From a point 0.5 mile upstream of backwaters of Kernersville Lake to Town of Kernersville Water Supply Dam		46.1 FW Acres	WS-IV;CA
3n	Chlorophyll a	Potential Standards Violation	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-29	Big Beaver Island Creek	From source to Dan River		15.2 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-(8)	DAN RIVER	From Big Creek to to a point 0.2 mile downstream of Town Fork Creek		25.9 FW Miles	WS-V
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Belews Lake-Dan River Watershed		0301010303	
⊙ 22-20-9	Lynn Branch (Lynn Creek)	From source to Snow Creek		3.1 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
⊙ 22-20-4	Raccoon Creek	From source to Snow Creek		3.4 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
⊙ 22-20	Snow Creek	From source to Dan River		18.9 FW Miles	C
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-27-9-(4)	West Belews Creek (West Belews Creek Arm of of Belews Lake below elevation 725)	From a point 0.4 mile downstream of Powerplant to Belews Creek		582.4 FW Acres	WS-IV
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-21	Wood Benton Branch	From source to Dan River		3.7 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2005	
Roanoke River Basin		Mayo River Watershed		0301010304	
⊙ 22-30-2-2	Crooked Creek (North Carolina portion)	From source to last crossing of North Carolina-Virginia State Line		8.5 FW Miles	C
1	Ecological/biological Integrity FishCom	Good-Fair Bioclassification	Aquatic Life	2007	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
⊙ 22-30-5	Hickory Creek	From source to Mayo River		4.0 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2006	
⊙ 22-30-2-2-2	Little Crooked Creek	From source to Crooked Creek		4.7 FW Miles	C
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2008	
⊙ 22-30-(1)	Mayo River	From North Carolina-Virginia State Line to a point 0.6 mile downstream of Hickory Creek		3.5 FW Miles	WS-V
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin			Mayo River Watershed		0301010304
⊙ 22-30-6-(2)	Pawpaw Creek	From a point 1.3 mile upstream of Rockingham County SR 1360 to Mayo R.		1.8 FW Miles	WS-IV
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
Roanoke River Basin			Matrimony Creek-Dan River Watershed		0301010305
⊙ 22-32-1	Brushy Creek (West Prong Jacobs Creek)	From source to Jacobs Creek		4.3 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2007	
⊙ 22-(31.5)a	DAN RIVER	From a point 0.7 mile upstream of Jacobs Creek to subbasin 03-02-02/03 boundary		4.8 FW Miles	WS-IV
4t	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	2008
4t	Turbidity	Data Inconclusive	Aquatic Life	2008	2002
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-(31.5)b	DAN RIVER	From 03-02-02 boundary to a point 0.8 mile downstream of Matrimony Creek		9.4 FW Miles	WS-IV
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008
4t	Turbidity	Data Inconclusive	Aquatic Life	2008	2002
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-(38.5)	DAN RIVER	From a point 0.8 mile downstream of Matrimony Creek to Mill Branch (Town of Eden water supply intake)		0.6 FW Miles	WS-IV;CA
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008
5	Turbidity	Standard Violation	Aquatic Life	2008	2008
⊙ 22-31	Hogans Creek	From source to Dan River		12.7 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-32-(3)	Jacobs Creek	From N.C. Hwy. 704 to Dan River		1.8 FW Miles	WS-IV
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-38	Matrimony Creek (North Carolina portion)	From source to Dan River		11.2 FW Miles	WS-IV
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Matrimony Creek-Dan River Watershed		0301010305	
⊙ 22-30-(10)	Mayo River	From dam at Mayodan Water Supply Intake to Dan River		2.4 FW Miles	C
1	Ecological/biological Integrity Benthos	Good-Fair Bioclassification	Aquatic Life	1999	
⊙ 22-34-(2)	Rock House Creek	From Rockingham County SR 2381 to Dan River		6.5 FW Miles	WS-IV
1	Ecological/biological Integrity Benthos	Good-Fair Bioclassification	Aquatic Life	2001	
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Lower Smith River Watershed		0301010308	
⊙ 22-40-(1)	Smith River	From North Carolina-Virginia State Line to a point 0.8 mile downstream of Rockingham County SR 1714 (Aiken Road)		2.8 FW Miles	WS-IV
5	Copper	Standard Violation	Aquatic Life	2008	2008
4s	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	1999	2008
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-40-(2.5)	Smith River	From a point 0.8 mile downstream of Rockingham County SR 1714 (Aiken Road) to Fieldcrest Mills Water Supply Intake		0.5 FW Miles	WS-IV;CA
5	Copper	Standard Violation	Aquatic Life	2008	2008
4s	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	1999	2008
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
Roanoke River Basin		Cascade Creek-Dan River Watershed		0301010309	
⊙ 22-48-4	Birch Fork	From source to Wolf Island Creek		8.4 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2007	
⊙ 22-(39)a	DAN RIVER (North Carolina portion)	From Mill Branch to NC/VA crossing downstream of Wolf Island Creek		13.8 FW Miles	C
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008
5	Turbidity	Standard Violation	Aquatic Life	2008	2008
⊙ 22-40-(3)	Smith River	From Fieldcrest Mills Water Supply Intake to Dan River		1.8 FW Miles	C
5	Copper	Standard Violation	Aquatic Life	2008	2008
4s	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	1999	2008
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Cascade Creek-Dan River Watershed		0301010309	
22-48	Wolf Island Creek	From source to Dan River		21.8 FW Miles	C
1	Ecological/biological Integrity FishCom	Excellent Bioclassification	Aquatic Life	2004	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin			Hogans Creek-Dan River Watershed		0301010401
Roanoke River Basin			Dan River Subbasin		03010104
Roanoke River Basin			Hogans Creek-Dan River Watershed		0301010401
22-54	Cane Creek	From North Carolina-Virginia State Line to Dan River		0.8 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
22-(39)b	DAN RIVER (North Carolina portion)	From NC/VA crossing downstream of Wolf Island Creek to last crossing of North Carolina-Virginia State Line		9.6 FW Miles	C
4t	Fecal Coliform (recreation)	Standard Violation	Recreation	2008	2008
5	Turbidity	Standard Violation	Aquatic Life	2008	2008
22-50	Hogans Creek	From source to Dan River		29.1 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
22-50-3	Jones Creek (Lake Wade)	From source to Hogans Creek		7.6 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
22-51	Moon Creek (Wildwood Lake)	From source to Dan River		17.0 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
22-52	Rattlesnake Creek	From source to Dan River		2.7 FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
Roanoke River Basin			Country Line Creek Watershed		0301010402
22-56-(1)	Country Line Creek	From source to a point 0.5 mile upstream of mouth of Nats Fork		10.5 FW Miles	WS-II;HQW
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
22-56-(3.7)	Country Line Creek	From dam at Farmer Lake to Dan River		24.5 FW Miles	C
1	Ecological/biological Integrity Benthos	Good Bioclassification	Aquatic Life	2004	
22-56-(3.5)a	Country Line Creek (Farmers Lake)	Upper reservoir- From a point 0.5 mile upstream of mouth Nats Fork to dam at Farmer Lake (Town of Yanceyville water supply intake located 1.8 mile upstream of N.C. Hwy. 62)		90.7 FW Acres	WS-II;HQW,CA
5	Chlorophyll a	Standard Violation	Aquatic Life	2008	2010
5	Turbidity	Standard Violation	Aquatic Life	2008	2010
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	

NC DWQ ROANOKE RIVER BASIN PLAN: 2010 USE SUPPORT 2011

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Country Line Creek Watershed		0301010402	
⊙ 22-56-(3.5)b	Country Line Creek (Farmers Lake)	Lower reservoir-From a point 0.5 mile upstream of mouth Nats Fork to dam at Farmer Lake (Town of Yanceyville water supply intake located 1.8 mile upstream of N.C. Hwy. 62)	271.1 FW Acres		WS-II;HQW,CA
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Water Supply	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
Roanoke River Basin		Hyco Lake Watershed		0301010405	
⊙ 22-58-1	Hyco Creek (North Hyco Creek)	From source to Hyco Lake, Hyco River	16.8 FW Miles		C
3a	Ecological/biological Integrity FishCom	Not Rated Bioclassification	Aquatic Life	2004	
⊙ 22-58-(0.5)	Hyco River, including Hyco Lake below elevation 410	From source in Hyco Lake to dam of Hyco Lake, including tributary arms below elevation 410	4,297.9 FW Acres		WS-V,B
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
⊙ 22-58-4-(3)	South Hyco Creek	From a point 0.6 mile downstream of Double Creek to Hyco Lake, Hyco River (City of Roxboro water supply intake)	0.7 FW Miles		WS-II;HQW,CA
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 22-58-4-(1.4)	South Hyco Creek (Lake Roxboro)	From backwaters of Lake Roxboro to dam at Lake Roxboro	493.6 FW Acres		WS-II,B;HQV
3n	Chlorophyll a	Potential Standards Violation	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
Roanoke River Basin		Hyco River Watershed		0301010406	
⊙ 22-58-(9.5)	Hyco River	From dam of Hyco Lake to North Carolina-Virginia State Line, including all portions in North Carolina	6.8 FW Miles		C
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
⊙ 22-58-12-6a	Marlowe Creek	From source to Mitchell Creek	6.6 FW Miles		C
5	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	2004	1998

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin			Hyco River Watershed		0301010406

22-58-12-6b	Marlowe Creek	From Mithcell Creek to Storys Creek	4.5	FW Miles	C
5	Copper	Standard Violation	Aquatic Life	2008	2008
1	Ecological/biological Integrity Benthos	Good-Fair Bioclassification	Aquatic Life	2004	
1	Ecological/biological Integrity FishCom	Good-Fair Bioclassification	Aquatic Life	2004	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
5	Zinc	Standard Violation	Aquatic Life	2008	2008
22-58-15-(3.5)	Mayo Creek (Maho Creek)	From dam of Mayo Reservoir to North Carolina-Virginia State Line	0.5	FW Miles	C
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
22-58-15-(0.5)	Mayo Creek (Maho Creek) (Mayo Reservoir)	From source to dam of Mayo Reservoir	2,613.8	FW Acres	WS-V
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
22-58-12-(1.5)	Storys Creek [Roxboro City Lake (Lake Issac Walton)]	From a point 0.9 mile downstream of N.C. Hwy. 57 to Roxboro City Lake Dam	189.5	FW Acres	WS-II;HQW,CA
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	

Roanoke River Basin	Aarons Creek-Dan River Watershed	0301010407
----------------------------	---	-------------------

22-59	Aarons Creek	From source to North Carolina-Virginia State Line	8.6	FW Miles	C
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	

NC DWQ ROANOKE RIVER BASIN PLAN: 2010 USE SUPPORT 2011

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Upper Lake Gaston-Roanoke River Watershed		0301010602	
Roanoke River Basin		Lake Gaston-Roanoke River Subbasin		03010106	
Roanoke River Basin		Upper Lake Gaston-Roanoke River Watershed		0301010602	
⊙ 23-10-2	Newmans Creek (Little Deep Creek)	From source to Smith Creek		6.1 FW Miles	C
5	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	2004	2008
⊙ 23-10a	Smith Creek	From source to Cabin Branch		6.1 FW Miles	C
4s	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	2004	2002
5	Low Dissolved Oxygen	Standard Violation	Aquatic Life	2008	1998
⊙ 23-10b	Smith Creek	From Cabin Branch to SR1208		1.6 FW Miles	C
1	Ecological/biological Integrity Benthos	Good-Fair Bioclassification	Aquatic Life	2004	
5	Low Dissolved Oxygen	Standard Violation	Aquatic Life	2008	1998
⊙ 23-10c	Smith Creek	From SR1208 to North Carolina-Virginia State Line		3.0 FW Miles	C
4s	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	2004	
4s	Ecological/biological Integrity FishCom	Fair Bioclassification	Aquatic Life	2004	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
5	Low Dissolved Oxygen	Standard Violation	Aquatic Life	2008	1998
⊙ 23-10-3-2	Terrapin Creek	From source to Blue Mud Creek		5.0 FW Miles	C
3a	Ecological/biological Integrity Benthos	Not Rated Bioclassification	Aquatic Life	2007	
Roanoke River Basin		Middle Lake Gaston-Roanoke River Watershed		0301010603	
⊙ 23-14	Jordan Creek	From source to Lake Gaston, Roanoke River		2.6 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2006	
⊙ 23-13	Sixpound Creek	From source to Lake Gaston, Roanoke River		6.3 FW Miles	C
1	Ecological/biological Integrity Benthos	Good-Fair Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Lower Lake Gaston-Roanoke River Watershed		0301010604	
⊙ 23-24-(1)	Deep Creek	From source to a point 0.5 mile upstream of mouth		11.6 FW Miles	WS-IV
1	Ecological/biological Integrity Benthos	Natural Bioclassification	Aquatic Life	2004	
1	Ecological/biological Integrity FishCom	Good Bioclassification	Aquatic Life	2004	
⊙ 23-19	Little Stonehouse Creek	From source to Lake Gaston, Roanoke River		2.8 FW Miles	C
1	Ecological/biological Integrity Benthos	Not Impaired Bioclassification	Aquatic Life	2006	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Lower Lake Gaston-Roanoke River Watershed		0301010604	

⊙ **23-(22.5)** **ROANOKE RIVER**
(Lake Gaston below normal full power pool elevation 200 MSL and Roanoke Rapids Lake below normal full power pool elevation 132 feet MSL)

From a line across Lake Gaston 0.5 mile upstream of Lake Gaston Dam to Roanoke Rapids Dam

4,185.0 FW Acres **WS-IV,B;CA**

3t	Aquatic Weeds	Data Inconclusive	Aquatic Life	2000	2000
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	

⊙ **23-(12)** **ROANOKE RIVER**
(Lake Gaston below normal full power pool elevation 200 MSL)

From North Carolina-Virginia State Line to a line across Lake Gaston following the Warren-Northampton County Line

7,964.8 FW Acres **WS-V,B**

1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
----------	--------------------------------------	----------------------	--------------	------	--

⊙ **23-(20.2)** **ROANOKE RIVER**
(Lake Gaston below normal full power pool elevation 200 MSL)

From a line across Lake Gaston following the Warren-Northampton County Line to a line across Lake Gaston 0.5 mile upstream of Lake Gaston Dam

3,974.4 FW Acres **WS-IV,B**

1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Quankey Creek-Roanoke River Watershed		0301010701	
Roanoke River Basin		Roanoke River Subbasin		03010107	
Roanoke River Basin		Quankey Creek-Roanoke River Watershed		0301010701	
23-29	Chockoyotte Creek	From source to Roanoke River		10.6 FW Miles	C
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
3a	Ecological/biological Integrity FishCom	Not Rated Bioclassification	Aquatic Life	2004	
23-30-1	Little Quankey Creek	From source to Quankey Creek		9.5 FW Miles	C
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
23-30a	Quankey Creek	From source to Little Quankey Creek		16.0 FW Miles	C
1	Ecological/biological Integrity Benthos	Natural Bioclassification	Aquatic Life	2004	
23-30b	Quankey Creek	From Little Quankey Creek to Roanoke River		3.4 FW Miles	C
5	Ecological/biological Integrity Benthos	Fair Bioclassification	Aquatic Life	1999	1998
23-(25.5)	ROANOKE RIVER	From a point 0.6 mile upstream of N.C. Hwy. 48 bridge to a line across river 50 feet downstream of N.C. Hwy. 48 (City of Roanoke Rapids, Town of Weldon water supply intakes)		1.7 FW Miles	WS-IV;CA
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
1	Water Quality Standards Water Supply	No Criteria Exceeded	Water Supply	2008	
23-(26)a	ROANOKE RIVER	From a line across the river 50 ft downstream of NC Hwy 48 bridge to the confluence of Sandy Run Cr at the Bertie Northampton Halifax Co. line		50.1 FW Miles	C
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
Roanoke River Basin		Conoconnara Swamp-Roanoke River Watershed		0301010702	
23-33	Conoconnara Swamp	From source to Roanoke River		17.7 FW Miles	C
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Kehukee Swamp-Roanoke River Watershed		0301010703	
23-42	Kehukee Swamp (White Millpond)	From source to Roanoke River		10.6 FW Miles	C
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin		Kehukee Swamp-Roanoke River Watershed		0301010703	
⊙ 23-(26)b1	ROANOKE RIVER	From the confluence of Sandy Run Cr at the Bertie/Northampton/Halifax Co. line to subbasin 8/9 boundary	24.8 FW Miles	C	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
Roanoke River Basin		Sweetwater Creek Watershed		0301010704	
⊙ 23-50-3	Hardison Mill Creek	From source to Sweetwater Creek	19.9 FW Miles	C	
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Conoho Creek-Roanoke River Watershed		0301010705	
⊙ 23-49a	Conoho Creek	From source to Martin Co 1417 below Beaverdam Cr	24.5 FW Miles	C	
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
⊙ 23-49b	Conoho Creek	From Martin Co 1417 to Roanoke River	7.0 FW Miles	C	
1	Ecological/biological Integrity Benthos	Natural Bioclassification	Aquatic Life	2004	
⊙ 23-(26)b2	ROANOKE RIVER	From subbasin 8/9 boundary to Hwy 17 Bridge in Williamston	28.9 FW Miles	C	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
⊙ 23-(26)b3	ROANOKE RIVER	From Hwy 17 bridge at Williamston to the 18 mile marker at Jamesville	17.8 FW Miles	C	
5	Low Dissolved Oxygen	Standard Violation	Aquatic Life	2006	2008
Roanoke River Basin		Headwaters Cashie River Watershed		0301010707	
⊙ 24-2-(1)a	Cashie River	From source to Bertie County SR 1225	15.2 FW Miles	C;Sw	
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	
Roanoke River Basin		Outlet Cashie River Watershed		0301010708	
⊙ 24-2-(1)b	Cashie River	From Bertie County SR 1225 to a point 1 mile upstream from Bertie Co. SR 1500	30.1 FW Miles	C;Sw	
1	Ecological/biological Integrity Benthos	Natural Bioclassification	Aquatic Life	2004	
⊙ 24-2-6	Hoggard Mill Creek	From source to Cashie River	7.4 FW Miles	C;Sw	
1	Ecological/biological Integrity Benthos	Moderate Bioclassification	Aquatic Life	2004	
⊙ 24-2-7	Roquist Creek	From source to Cashie River	26.3 FW Miles	C;Sw	
1	Ecological/biological Integrity Benthos	Natural Bioclassification	Aquatic Life	2004	
Roanoke River Basin		Plymouth-Roanoke River Watershed		0301010709	

NC 2010 Integrated Report

All 13,123 Waters in NC are in Category 5-303(d) List for Mercury due to statewide fish consumption advice for several fish species

AU_Number	AU_Name	AU_Description	LengthArea	AU_Units	Classification
Category	Parameter	Reason for Rating	Use Category	Collection Year	303(d)year
Roanoke River Basin			Plymouth-Roanoke River Watershed		0301010709

⊙ **23-(53)** **ROANOKE RIVER** From 18 mile marker at Jamesville to Albemarle Sound (Batchelor Bay) **18.3 FW Miles** C;Sw

4t	Dioxin	Standard Violation	Fish Consumption	2008	2000
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
1	Water Quality Standards Aquatic Life	No Criteria Exceeded	Aquatic Life	2008	

⊙ **23-55** **Welch Creek** From source to Roanoke River **13.3 FW Miles** C;Sw

4t	Dioxin	Standard Violation	Fish Consumption	1996	2000
1	Fecal Coliform (recreation)	No Criteria Exceeded	Recreation	2008	
5	Low pH	Standard Violation	Aquatic Life	2008	2002

2010
Use Assessment Methodology
EPA Approved August 31, 2010

Table of Contents

Purpose.....	3
Assessment Units and Water Quality Classifications	3
Data Window/Assessment Period	4
Data Availability and Quality	4
Use Support Categories and Water Quality Standards	4
Aquatic Life Assessment Methodology	4
Numerical Water Quality Standards	4
Dissolved Oxygen (DO) Standards	5
Freshwater Dissolved Oxygen (DO) Assessment (Class C, B, WS).....	5
Saltwater Dissolved Oxygen (DO) Assessment (Class SC, SB, SA).....	5
Trout Water Dissolved Oxygen (DO) Assessment (Supplemental Class Tr).....	5
Swamp Water Dissolved Oxygen (DO) Assessment (Supplemental Class Sw)	5
pH.....	6
pH Standards	6
Low pH Assessment (Class C, SC, B, SB, SA, WS).....	6
High pH Assessment (Class C, SC, B, SB, SA, WS).....	6
Swamp Water Low pH Assessment (Supplemental Class Sw).....	7
Temperature Use Assessment.....	7
Temperature Standards	7
Temperature Assessment.....	7
Temperature Screening Criteria for Trout Waters (Supplemental Class Tr)	8
Assessment of Extreme Temperature Conditions	8
Chlorophyll <i>a</i>	8
Chlorophyll <i>a</i> Standard	8
Chlorophyll <i>a</i> Standards Assessment	8
Toxic Substances and Action Levels Metals	9
Toxic Substances Numerical Standards	9
Metals Action Level Standards.....	9
Toxic Substances and Action Level Metals Assessment.....	9
Turbidity	10
Turbidity Standards	10
Turbidity Assessment.....	10
Ecological/Biological Integrity.....	11
Aquatic Life Narrative Standards.....	11
Aquatic Life Assessment	11
Recreation Assessment Methodology.....	11
Pathogen Indicator Standards	12
Fecal Coliform Bacteria Assessment Criteria.....	12
Fecal Coliform Bacteria Screening Assessment.....	12
Enterococci Assessment Criteria.....	12
Enterococcus Screening Assessment.....	12
Advisory Posting Assessment	12
Shellfish Harvesting Assessment Methodology	13
Shellfish Harvesting Standards	13
Fecal Coliform Bacteria Assessment Criteria.....	13
DEH Shellfish Sanitation Growing Area Classification Assessment.....	13
Water Supply Assessment Methodology.....	13

Water Supply Standards.....	13
Water Supply Assessment.....	13
Fish Consumption Assessment Methodology	14
Polychlorinated biphenyls (PCBs) Assessment Criteria	14
Dioxin Assessment Criteria	14
Mercury Assessment Criteria	14

Purpose

Section 303(d) of the federal Clean Water Act (CWA) which Congress enacted in 1972 requires States, Territories and authorized Tribes to identify and establish a priority ranking for waterbodies for which technology-based effluent limitations required by section 301 are not stringent enough to attain and maintain applicable water quality standards, establish total maximum daily loads (TMDLs) for the pollutants causing impairment in those waterbodies, and submit, from time to time, the list of impaired waterbodies and TMDLs to the U.S. Environmental Protection Agency (EPA). Current federal rules require states to submit 303(d) lists biennially, by April 1st of every even numbered year. The “303(d) list” is technically considered the impaired waters listed as Category 5, requiring a TMDL. EPA is required to approve or disapprove the state-developed §303(d) list within 30 days. For each water quality limited segment impaired by a pollutant and identified in the §303(d) list, a Total Maximum Daily Load (TMDL) must be developed.

Assessment Units and Water Quality Classifications

Water quality assessments are based on water quality classifications as well as data availability. Water quality classifications are associated with a stream reach or area that is described in the schedule of classifications. Reaches vary in length or area and are sometimes split into smaller units to represent application of water quality data. Classifications are represented by a series of numbers called index numbers, 27-33-43-(1), as an example. Water quality assessments are applied to assessment units or AUs. AUs are, for the most part, the same as index numbers. When an AU is subdivided because of data applicability a letter is added to indicate this smaller unit. For example, if Index number 27-33-43-(1) (12 miles in length) is divided into three different segments because of three different available data types the new segments would be 27-33-43-(1)a, 27-33-43-(1)b and 27-33-43-(1)c. The combined mileage of the AUs would be 12 miles.

Decisions on the length or area to apply data to are based on the data type, waterbody characteristics, stations indicating similar water quality, watershed information and landmarks on which to base descriptions. The AUs where water quality concerns are evident are used as markers. Solutions to water quality concerns, including TMDLs, typically encompass entire watersheds.

Data Window/Assessment Period

The data window for the 2010 Water Quality Use Assessment (305(b) and 303(d) Integrated Reporting) includes data collected in calendar years 2004 through 2008 (five years). Some AUs may have biological data collected earlier for waters that have not been resampled during this data window or where the current impairment is based on that sample. The data collection year is noted for each AU.

Data Availability and Quality

Data are collected by various state and federal agencies. NC Department of Environment and Natural Resources (NCDENR) Division of Water Quality (DWQ) collects most of the data used for water quality assessments. There are significant data sets collected by NCDENR Division of Environmental Health (DEH) for use in coastal water quality assessment. The United States Geological Survey (USGS) also provides data in several AUs. Local governments and environmental groups as well as industry, municipal and university coalitions also provide data. Submitted data sets must include an approved Quality Assurance Project Plan (QAPP) or other documentation to assure that the data were collected in a manner consistent with agency data. A standing solicitation for data is maintained on the DWQ website. DWQ evaluates all data and information submitted.

Use Support Categories and Water Quality Standards

There are numerical and narrative water quality standards that are in place to protect the various best uses of North Carolina waters. Best uses include aquatic life or biological integrity, recreation or swimming, fish consumption, shellfish harvesting and water supply. Water quality assessments are based on the standards and data availability for the applicable use support category- aquatic life, recreation etc. Dissolved oxygen standards are used to assess aquatic life and pathogen indicators are used to assess recreation for example. Standards assessment criteria have been developed for each parameter assessed. The standards assessment criteria are used to make water quality assessments- not the standards themselves. While the standards assessment criteria are based on the standards they are different in that a frequency term is included. The details of how each standard is assessed are discussed in the following sections.

Aquatic Life Assessment Methodology

Numerical Water Quality Standards

The aquatic life numerical water quality standards are assessed using a 10% exceedance of the standard criterion. These assessments use ambient monitoring data from the five year assessment period (2004-2008). If no aquatic life numerical water quality standards exceed the 10% criterion then the AU is Supporting aquatic life water quality standards. This AU/multiple-parameters assessment is a Category 1 listing not requiring a TMDL. If greater than 10% of the

samples exceed the numerical standard and there are at least 10 samples, then the AU is Impaired for that parameter. The AU/parameter assessment is listed in Category 5, requiring a TMDL. If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL. The NC DWQ "Redbook" contains the complete descriptions of water quality standards and surface water classifications [15a NCAC 02B .0200 - .0300]

Dissolved Oxygen (DO) Standards

Freshwater dissolved oxygen: not less than 6.0 mg/l for trout waters; for non-trout waters, not less than a daily average of 5.0 mg/l with a minimum instantaneous value of not less than 4.0 mg/l; swamp waters, lake coves or backwaters, and lake bottom waters may have lower values if caused by natural conditions.

Salt water dissolved oxygen: not less than 5.0 mg/l, except that swamp waters, poorly flushed tidally influenced streams or embayments, or estuarine bottom waters may have lower values if caused by natural conditions.

Freshwater Dissolved Oxygen (DO) Assessment (Class C, B, WS)

A fresh non-swamp water AU was assessed as Impaired for aquatic life when greater than 10% of samples were below 4 mg/l for instantaneous samples (monthly) or when greater than 10% of samples are below a daily average of 5mg/l. A minimum of 10 samples was needed to rate the water as Impaired.

Saltwater Dissolved Oxygen (DO) Assessment (Class SC, SB, SA)

A saline/estuarine non-swamp water AU was assessed as Impaired for aquatic life when greater than 10% of samples were below 5 mg/l. A minimum of 10 samples was needed to rate the water as Impaired.

Trout Water Dissolved Oxygen (DO) Assessment (Supplemental Class Tr)

A supplemental classified Trout water AU was assessed as Impaired for aquatic life when greater than 10% of samples were below 6 mg/l. A minimum of 10 samples was needed to rate the water as Impaired.

Swamp Water Dissolved Oxygen (DO) Assessment (Supplemental Class Sw)

A supplemental classified swamp (Sw) AU was Not Rated for aquatic life when greater than 10% of samples were below 4 mg/l (5 mg/l for salt) for instantaneous samples (monthly) or when greater than 10% of samples were below a daily average of 5 mg/l (freshwater only). There is not a numerical standard for these waterbodies and natural background conditions cannot be determined. This is a category 3a listing not requiring a TMDL.

A swamp like AU (not classified Sw) was Not Rated for aquatic life when greater than 10% of samples were below 4 mg/l (5 mg/l for salt) for instantaneous samples

(monthly) or when greater than 10% of samples were below a daily average of 5mg/l (freshwater only) and when greater than 10% of samples were below a pH of 6.0 (SU) for freshwater or 6.8 (SU) for saltwater. Geographic location, biological data, tributary classifications, discharges and land use were considered when assigning use support ratings to waters considered to be swamp like or receiving significant swamp water input.

pH

pH Standards

Freshwater pH: shall be normal for the waters in the area, which generally shall range between 6.0 and 9.0 except that swamp waters may have a pH as low as 4.3 if it is the result of natural conditions;

Saltwater pH: shall be normal for the waters in the area, which generally shall range between 6.8 and 8.5 except that swamp waters may have a pH as low as 4.3 if it is the result of natural conditions;

Low pH Assessment (Class C, SC, B, SB, SA, WS)

A non-swamp water AU was assessed as Impaired for aquatic life when greater than 10% of samples were below a pH of 6.0 (SU) for freshwater or 6.8 (SU) for saltwater.

A swamp like AU (not classified Sw) was Not Rated for aquatic life when greater than 10% of samples were below a pH of 6.0 (SU) for freshwater or 6.8 (SU) for saltwater or when greater than 10% of samples were below a dissolved oxygen of 4 mg/l (5 mg/l for salt) for instantaneous samples (monthly) or when greater than 10% of samples were below a daily average of 5mg/l (freshwater only) Geographic location, biological data, tributary classifications, discharges and land use were considered when making use support determinations on waters considered to be swamp like or receiving significant swamp water input.

High pH Assessment (Class C, SC, B, SB, SA, WS)

An AU was assessed as Impaired for aquatic life when greater than 10% of samples were greater than a pH of 9 (SU) for freshwater or 8.5 (SU) for saltwater. A minimum of 10 samples was needed to rate the water as Impaired. This is a Category 5 listing requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

Swamp Water Low pH Assessment (Supplemental Class Sw)

A supplemental classified swamp (Sw) AU was assessed as Impaired when greater than 10% of samples were below 4.3 (SU). A minimum of 10 samples was needed to rate the water as Impaired. This is a Category 5 listing requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

Temperature Use Assessment

Temperature Standards

For freshwaters- Temperature: not to exceed 2.8°C (5.04°F) above the natural water temperature, and in no case to exceed 29°C (84.2°F) for mountain and upper piedmont waters and 32°C (89.6°F) for lower piedmont and coastal plain waters. The temperature for trout waters shall not be increased by more than 0.5°C (0.9°F) due to the discharge of heated liquids, but in no case to exceed 20°C (68°F).

Lower piedmont and coastal plain waters mean those waters of the Catawba River Basin below Lookout Shoals Dam; the Yadkin River Basin below the junction of the Forsyth, Yadkin, and Davie County lines; and all of the waters of Cape Fear, Lumber, Roanoke, Neuse, Tar-Pamlico, Chowan, Pasquotank, and White Oak River Basins; except tidal salt waters which are assigned S classifications.

Mountain and upper piedmont waters mean all of the waters of the Hiwassee; Little Tennessee, including the Savannah River drainage area; French Broad; Broad; New; and Watauga River Basins; and those portions of the Catawba River Basin above Lookout Shoals Dam and the Yadkin River Basin above the junction of the Forsyth, Yadkin, and Davie County lines.

For saltwaters- Temperature: shall not be increased above the natural water temperature by more than 0.8°C (1.44°F) during the months of June, July, and August nor more than 2.2°C (3.96°F) during other months and in no cases to exceed 32°C (89.6°F) due to the discharge of heated liquids.

Temperature Assessment

A mountain or upper piedmont AU was assessed as Impaired for aquatic life when greater than 10% of samples were greater than 29°C. A minimum of 10 samples was needed to rate the water as Impaired.

A lower piedmont or coastal plain stream AU was assessed as Impaired for aquatic life when greater than 10% of samples were greater than 32°C. A minimum of 10 samples was needed to rate the water as Impaired.

If the 10% criterion was exceeded and fewer than 10 samples were collected the water was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

Temperature Screening Criteria for Trout Waters (Supplemental Class Tr)

A supplemental classified trout water (Tr) AU was Not Rated for aquatic life when greater than 10% of samples were greater than 20°C. The presence of heated discharges was not determined. This is a Category 3a listing not requiring a TMDL.

Assessment of Extreme Temperature Conditions

A waterbody that exceeds the above criteria may be Not Rated for aquatic life because of meteorological conditions that occur on a regular basis. These conditions must be documented and reassessment will occur after more normal conditions return. This is a Category 3a listing not requiring a TMDL. Examples of extreme conditions may include extreme drought, reservoir drawdown, hurricane impacts and flooding, dam failure, and saltwater encroachment. Other extreme conditions may be documented as needed for future assessments

Chlorophyll *a*

Chlorophyll *a* Standard

Chlorophyll *a* (corrected): not greater than 40 µg/l in sounds, estuaries, and other waters subject to growths of macroscopic or microscopic vegetation.

Other waters subject to growths are interpreted by DWQ to include dam backwaters, lakes and reservoirs.

Chlorophyll *a* Standards Assessment

An AU was assessed as Impaired for aquatic life when greater than 10% of samples were greater than 40 µg/l. A minimum of 10 samples was needed to rate the water as Impaired. This is a Category 5 listing requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. Some reservoirs in North Carolina are sampled fewer than 10 times during the assessment period. These data are used to document eutrophication issues. Reservoirs are targeted for increased monitoring to determine if there are standards violations using the above methodology. This is a Category 3a listing not requiring a TMDL.

Toxic Substances and Action Levels Metals

Toxic Substances Numerical Standards

Refer to the NC DWQ "Redbook" for complete text of standards

Arsenic: 50 ug/l

Beryllium: 6.5 ug/l;

Cadmium: 0.4 ug/l for trout waters and 2.0 ug/l for non-trout waters;

Chlorine, total residual: 17 ug/l;

Chromium, total recoverable: 50 ug/l;

Cyanide: 5.0 ug/l

Fluorides: 1.8 mg/l;

Lead, total recoverable: 25 ug/l;

Mercury (assessed in fish consumption category)

Nickel: 88 ug/l; 8.3 ug/l

Chlorides: 230mg/l; (note this is an action level standard)

Metals Action Level Standards

Action Level Copper: 7 ug/l FW or 3 ug/l SW

Action Level Silver: 0.06 ug/l;

Action Level Zinc: 50 ug/l;

Toxic Substances and Action Level Metals Assessment

An AU was assessed as Impaired for aquatic life when greater than 10% of samples were greater than the above standards or action level standards. A minimum of 10 samples was needed to rate the water as Impaired. These are Category 5 listings requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

The action level standard for Iron was not assessed during this assessment period because the standard is being reevaluated and the Iron exceedances of the Action Level have been shown to be a natural condition.

Action levels are used for permitting purposes and are not used as the only information to assess aquatic life uses. Copper and Zinc may be indicators of potential impacts to aquatic life. DWQ will review Copper and Zinc assessments that result in Category 5 listings. The review will be used to determine if the Category 5 listing is appropriate. The following criteria will be used to determine if a review is warranted.

1. A collocated Good, Excellent, Natural or Not Impaired biological rating or

2. A collocated Good-Fair, Moderate or Not Rated biological rating and less than 25% of Copper or Zinc samples exceed the evaluation level.
3. There are no biological data available and less than 25% of Copper or Zinc samples exceed the evaluation level.

The Water Quality Assessment Team will evaluate and integrate the following lines of watershed information to determine if a Category 5 listing for Copper and/or Zinc is warranted.

- 1- Analysis of duration, frequency and magnitude of exceedances.
- 2- Historical data and trends for the parameter of interest.
- 3- Detailed assessment of all available biological data.
- 4- Qualitative aquatic habitat information.
- 5- Natural or background conditions assessment including current imagery.
- 6- Sample quality (note that Zinc samples can be easily contaminated)
- 7- Waterbody classifications and other designated uses.
- 8- Exceedances of other likely associated metals.
- 9- Biological data in nearby Assessment Units.
- 10- Potential Sources of metals
- 11- Site specific hardness

After review the Assessment team will determine if the AU/parameter assessment is more appropriately listed in a Category other than 5. Each reviewed assessment will require documented justification for a final Integrate Report category other than Category 5.

Turbidity

Turbidity Standards

Turbidity: the turbidity in the receiving water shall not exceed 50 Nephelometric Turbidity Units (NTU) in streams not designated as trout waters and 10 NTU in streams, lakes or reservoirs designated as trout waters; for lakes and reservoirs not designated as trout waters, the turbidity shall not exceed 25 NTU; if turbidity exceeds these levels due to natural background conditions, the existing turbidity level cannot be increased.

Turbidity Assessment

An AU was assessed as Impaired for aquatic life when greater than 10% of samples were greater than 50 NTU or 10 NTU for Tr waters or 25 NTU for lakes, reservoirs and estuarine waters. A minimum of 10 samples was needed to rate the water as Impaired. This is a Category 5 listing requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

Ecological/Biological Integrity

Aquatic Life Narrative Standards

The aquatic life narrative water quality standard is assessed using a biological integrity index criterion (or bioclassification). Biological integrity means the ability of an aquatic ecosystem to support and maintain a balanced and indigenous community of organisms having species composition, diversity, population densities and functional organization similar to that of reference conditions. Waters shall be suitable for aquatic life propagation and maintenance of biological integrity, wildlife, secondary recreation, and agriculture. Sources of water pollution which preclude any of these uses on either a short-term or long-term basis shall be considered to be violating a water quality standard.

Aquatic Life Assessment

An AU was assessed as Impaired for aquatic life when a fish or benthic macroinvertebrate community sample received a bioclassification of Severe, Poor or Fair and there were no other Aquatic Life standards violations. This is a Category 5 listing requiring a TMDL.

An AU was assessed as Impaired for aquatic life when a fish or benthic macroinvertebrate community sample received a bioclassification of Severe, Poor or Fair and there were other Aquatic Life numeric standards violations. This is a Category 4s listing requiring a TMDL for the identified aquatic life numerical standards violation (Category 5 or 4t listing) impairing the ecological/biological integrity of the waterbody.

An AU was assessed as Impaired for aquatic life when a fish or benthic macroinvertebrate community sample received a bioclassification of Severe, Poor or Fair and an approved TMDL for an aquatic life numerical water quality standard has been completely implemented. This is a Category 5s listing requiring a TMDL.

Recreation Assessment Methodology

Recreation standards were assessed using fecal coliform bacteria data collected at DWQ ambient stations and special study sites and enterococci data collected at DEH Recreational Monitoring sites in coastal waters. Screening criteria were used to assess areas for potential standards violations. DEH advisory postings were also used for recreation assessments as well. The following criteria were used to assess waters for recreation.

Pathogen Indicator Standards

Organisms of coliform group: fecal coliforms not to exceed geometric mean of 200/100 ml (MF count) based on at least five consecutive samples examined during any 30-day period and not to exceed 400/100 ml in more than 20 percent of the samples examined during such period.

Enterococcus, including *Enterococcus faecalis*, *Enterococcus faecium*, *Enterococcus avium* and *Enterococcus gallinarium*: not to exceed a geometric mean of 35 enterococci per 100 ml based upon a minimum of five samples within any consecutive 30 days.

Fecal Coliform Bacteria Assessment Criteria

An AU was assessed as Impaired when the geometric mean was greater than 200 colonies/100ml or greater than 20% of the samples were higher than 400 colonies/100ml. At least 5 samples must have been collected within the same 30-day period. This is a Category 5 listing requiring a TMDL.

Fecal Coliform Bacteria Screening Assessment

An AU was Not Rated when the geometric mean was greater than 200 colonies/100ml or greater than 20% of the samples were higher than 400 colonies/100ml. Samples were not collected in the same 30-day period. This is a Category 3a listing not requiring a TMDL. These AUs are prioritized for resampling 5 times in 30 days based on classification and available resources. Data are reviewed yearly for prioritization.

Enterococci Assessment Criteria

An AU was assessed as Impaired when the geometric mean was greater than 35 colonies/100ml. At least 5 samples must have been collected within the same 30-day period. This is a Category 5 listing requiring a TMDL.

Enterococcus Screening Assessment

An AU was Not Rated when the geometric mean was greater than 35 colonies/100ml. Samples were not collected in the same 30-day period. This is a Category 3a listing not requiring a TMDL.

Advisory Posting Assessment

An AU was assessed as Impaired when a swimming advisory was posted for greater than 61 days in any 5 year period (includes permanent postings). This is a Category 4cr listing not requiring a TMDL.

Shellfish Harvesting Assessment Methodology

Shellfish Harvesting standards were assessed using DEH growing area classifications. The following criteria were used to assess waters for shellfish harvesting.

Shellfish Harvesting Standards

Organisms of coliform group: fecal coliform group not to exceed a median MF of 14/100 ml and not more than 10% of the samples shall exceed an MF count of 43/100 ml in those areas most probably exposed to fecal contamination during the most unfavorable hydrographic and pollution conditions.

Fecal Coliform Bacteria Assessment Criteria

DEH fecal coliform data were not assessed to determine standards violations. Category 5 impairments were based on Growing Area Classifications alone.

DEH Shellfish Sanitation Growing Area Classification Assessment

An AU was assessed as Impaired when the DEH growing area classification was Prohibited or conditionally approved. This is a Category 5 listing requiring a TMDL.

Water Supply Assessment Methodology

Water Supply standards were assessed using data collected at DWQ ambient stations located in Class WSI-WSV waters. The following criteria were used to Impair waters for water supply. Category 5 listings were only made when Standards Assessment Criteria (SAC) were exceeded.

Water Supply Standards

Refer to Water Quality "Redbook" for complete text of standards

Barium: 1.0 mg/l;

Chloride: 250 mg/l;

Manganese: 200 ug/l; (not human health or aquatic life- not assessed)

Nickel: 25 ug/l;

Nitrate nitrogen: 10.0 mg/l;

2,4-D: 100 ug/l;

2,4,5-TP (Silvex): 10 ug/l;

Sulfates: 250 mg/l;

Water Supply Assessment

An AU was assessed as Impaired for water supply when greater than 10% of samples were greater than the above standards except for manganese. A minimum of 10 samples was needed to rate the water as Impaired. This is a Category 5 listing requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

Fish Consumption Assessment Methodology

Fish Consumption was assessed based on site-specific fish consumption advisories. The advisories were based on the NC Department of Health and Human Services (DHHS) consumption advisories developed using fish tissue data that exceed standards. The following criteria were used to Impair waters for fish consumption. Because of the statewide Mercury advice there were no use cases for Supporting fish consumption and therefore no overall Category 1 waters.

Polychlorinated biphenyls (PCBs) Assessment Criteria

An AU was assessed as Impaired when a site-specific advisory was posted for PCBs. This is a Category 5 listing requiring a TMDL.

Dioxin Assessment Criteria

An AU was assessed as Impaired when a site-specific advisory was posted for dioxins. This is a Category 5 listing requiring a TMDL.

Mercury Assessment Criteria

An AU was assessed as Impaired for fish consumption when greater than 10% of samples were greater than 0.012 $\mu\text{g/l}$. A minimum of 10 samples was needed to rate the water as Impaired. This is a Category 5 listing requiring a TMDL.

If the 10% criterion was exceeded and fewer than 10 samples were collected the AU was Not Rated and targeted for further sampling. This is a Category 3a listing not requiring a TMDL.

Statewide advice for Mercury in fish tissue was not assessed because it was not associated with a specific AU but was applied to all waters of the state. All AUs are considered Impaired and in Category 5 for the statewide Mercury fish consumption advice. Previous site specific listings for Mercury will no longer be listed in Category 5. DWQ continues to monitor mercury in fish tissue, and has identified specific locations where Mercury levels exceed 0.4mg/kg of fish tissue.